

Review

Perioperative nutrition: Recommendations from the ESPEN expert group



Dileep N. Lobo ^{a, b, *}, Luca Gianotti ^c, Alfred Adiamah ^a, Rocco Barazzoni ^d, Nicolaas E.P. Deutz ^e, Ketan Dhatariya ^f, Paul L. Greenhaff ^b, Michael Hiesmayr ^g, Dorte Hjort Jakobsen ^h, Stanislaw Klek ⁱ, Zeljko Krznaric ^j, Olle Ljungqvist ^{k, l}, Donald C. McMillan ^m, Katie E. Rollins ^a, Marina Panisic Sekeljic ⁿ, Richard J.E. Skipworth ^o, Zeno Stanga ^p, Audrey Stockley ^q, Ralph Stockley ^q, Arved Weimann ^r

^a Gastrointestinal Surgery, Nottingham Digestive Diseases Centre, National Institute for Health Research (NIHR) Nottingham Biomedical Research Centre, Nottingham University Hospitals NHS Trust and University of Nottingham, Queen's Medical Centre, Nottingham, NG7 2UH, UK

^b MRC Versus Arthritis Centre for Musculoskeletal Ageing Research, National Institute for Health Research (NIHR) Nottingham Biomedical Research Centre, School of Life Sciences, University of Nottingham, Queen's Medical Centre, Nottingham, NG7 2UH, UK

^c School of Medicine and Surgery, University of Milano-Bicocca, Department of Surgery, San Gerardo Hospital, Monza, Italy

^d Department of Medical, Surgical and Health Sciences, University of Trieste, Trieste, Italy

^e Center for Translational Research in Aging & Longevity, Department of Health & Kinesiology, Texas A&M University, College Station, TX, 77843-4253, USA

^f Department of Diabetes, Endocrinology and General Medicine, Elsie Bertram Diabetes Centre, Norfolk and Norwich University Hospitals NHS Foundation Trust and University of East Anglia, Colney Lane, Norwich, NR4 7UY, UK

^g Division of Cardio-Thoracic-Vascular Surgical Anaesthesia and Intensive Care Medicine, Medical University of Vienna, Vienna, Austria

^h Section of Surgical Pathophysiology 4074, Rigshospitalet, Copenhagen, Denmark

ⁱ General and Oncology Surgery Unit, Stanley Dudrick's Memorial Hospital, Skawina, Poland

^j University Hospital Centre Zagreb and Zagreb School of Medicine, University of Zagreb, Zagreb, Croatia

^k Faculty of Medicine and Health, School of Health and Medical Sciences, Department of Surgery, Örebro University, Örebro, Sweden

^l Department of Molecular Medicine and Surgery, Karolinska Institutet, Stockholm, Sweden

^m Academic Unit of Surgery, School of Medicine, University of Glasgow, Royal Infirmary, Glasgow, UK

ⁿ Military Medical Academy, Clinic for General Surgery, Department for Perioperative Nutrition, Crnostravska Street 17, Belgrade, Serbia

^o Clinical Surgery, University of Edinburgh, Royal Infirmary of Edinburgh, 51 Little France Crescent, Edinburgh, EH16 4SA, UK

^p Division of Diabetes, Endocrinology, Nutritional Medicine and Metabolism, Bern University Hospital and University of Bern, Bern, Switzerland

^q Patient Public Involvement Group, Nottingham University Hospitals NHS Trust, Queen's Medical Centre, Nottingham, NG7 2UH, UK

^r Klinik für Allgemein-, Viszeral- und Onkologische Chirurgie, Klinikum St. Georg gGmbH, Delitzscher Straße 141, 04129, Leipzig, Germany

ARTICLE INFO

Article history:

Received 24 March 2020

Accepted 24 March 2020

Keywords:

Perioperative nutrition

Malnutrition

Nutritional assessment

Nutritional intervention

Perioperative care

Sarcopenia

SUMMARY

Background & aims: Malnutrition has been recognized as a major risk factor for adverse postoperative outcomes. The ESPEN Symposium on perioperative nutrition was held in Nottingham, UK, on 14–15 October 2018 and the aims of this document were to highlight the scientific basis for the nutritional and metabolic management of surgical patients.

Methods: This paper represents the opinion of experts in this multidisciplinary field and those of a patient and caregiver, based on current evidence. It highlights the current state of the art.

Results: Surgical patients may present with varying degrees of malnutrition, sarcopenia, cachexia, obesity and myosteatosis. Preoperative optimization can help improve outcomes. Perioperative fluid therapy should aim at keeping the patient in as near zero fluid and electrolyte balance as possible. Similarly, glycemic control is especially important in those patients with poorly controlled diabetes, with a stepwise increase in the risk of infectious complications and mortality per increasing HbA1c. Immobilization can induce a decline in basal energy expenditure, reduced insulin sensitivity, anabolic resistance to protein nutrition and muscle strength, all of which impair clinical outcomes. There is a role for pharmaconutrition, pre-, pro- and syn-biotics, with the evidence being stronger in those undergoing surgery for gastrointestinal cancer.

* Corresponding author. Gastrointestinal Surgery, Nottingham Digestive Diseases Centre, Nottingham University Hospitals NHS Trust and University of Nottingham, Queen's Medical Centre, E Floor, West Block, Nottingham, NG7 2UH, UK. Fax: +44 115 8231160.

E-mail address: Dileep.Lobo@nottingham.ac.uk (D.N. Lobo).

Conclusions: Nutritional assessment of the surgical patient together with the appropriate interventions to restore the energy deficit, avoid weight loss, preserve the gut microbiome and improve functional performance are all necessary components of the nutritional, metabolic and functional conditioning of the surgical patient.

© 2020 Elsevier Ltd and European Society for Clinical Nutrition and Metabolism. All rights reserved.

1. Introduction

Major surgery evokes a catabolic response that results in inflammation, protein catabolism and nitrogen losses. This response is proportional to the magnitude of the procedure and can, in some instances, be detrimental to the patient, especially when there is pre-existing malnutrition. Traditional perioperative care has involved measures that starve the patient for prolonged periods of time, stress the patient with interventions that amplify the catabolic response and drown the patient with salt and water overload. However, over the past two decades, there has been a paradigm shift in perioperative care, with periods of starvation being reduced drastically, introduction of measures to reduce surgical stress and protein catabolism, and avoiding salt and water overload. The aim of modern perioperative care is to attenuate loss of or aid functional recovery in an accelerated manner by promoting return of gastrointestinal function, feeding the patient early, providing adequate pain relief, and encouraging early mobilization. These measures result in reduced complications, early discharge from hospital without increasing readmission rates, and better functional recovery.

The European Society for Clinical Nutrition and Metabolism (ESPEN) has published updated evidence-based guidelines on perioperative nutrition recently that help aid the nutritional care of the surgical patient [1]. In further support of these guidelines, an ESPEN expert group met for a Perioperative Nutrition Symposium in Nottingham, UK on October 14 and 15, 2018. The group examined the causes and consequences of preoperative malnutrition, reviewed currently available treatment approaches in the pre- and postoperative periods, and analyzed the rationale on which clinicians could take actions that facilitate optimal nutritional and metabolic care in perioperative practice. The content of this position paper is based on presentations and discussions at the Nottingham meeting along with a subsequent update of the literature.

2. Historical note

Our understanding of the concept of clinical nutrition and the science of human nutrition has evolved significantly over the last two decades. The role of nutrition in surgery has encompassed measures to recognize, identify and intervene in those preoperative patients who are at risk of malnutrition with appreciable impact on postoperative outcomes in those adequately nutritionally prehabilitated. However, it would be incorrect to consider clinical nutrition as an entirely new concept [2–4]. Ancient Egyptians were the first to be credited with descriptions befitting enteral nutritional as identified in the Ebers papyrus (c 1550 BC) [4] and feeding via the oropharyngeal and nasopharyngeal routes are from then on described throughout the antiquated medical literature. For instance, Capivaccus in the 16th century, Aquapendente in the 17th century [2,4] and the 19th century physician Dukes [5] employed these routes of nutritional delivery to treat all manner of ailments including mania, diphtheria and croup.

The recognition of nutritional deficiency as a cause of illness was first presented by James Lind, a fellow of the Royal College of Physicians of Edinburgh who established the superiority of citrus fruits above all other 'remedies' in his treatise on scurvy published in 1753 [6]. The identification, characterization and synthesis of essential vitamins and minerals during the earlier part of the 20th century [7], allowing their use in the treatment of nutritional deficiency-related diseases such as scurvy, pellagra, rickets, and nutritional anemias [7].

The adverse effect of weight loss on surgical outcome was documented over 80 years ago when Hiram Studley showed that, in patients undergoing surgery for perforated duodenal ulcer, postoperative mortality was 10 times greater in those who had lost more than 20% of their body weight preoperatively when compared with those who had lost less [8]. This observation generated much of the ensuing work to define the role of malnutrition, nutritional deficiencies, and perioperative nutrition in surgery.

3. The malnourished surgical patient

The definition of a malnourished patient is the subject of ongoing discussion. In the last decade there have been considerable efforts to rationalize various definitions generally, and in the cancer patient for whom surgery is commonly the primary modality for cure. The starting point for much of this work was the international consensus of 2011 [9]. In this publication, cancer cachexia was defined as "a multifactorial syndrome defined by an ongoing loss of skeletal muscle mass (with or without loss of fat mass) that cannot be fully reversed by conventional nutritional support and leads to progressive functional impairment." There was a recognition of the role of the systemic inflammatory response in the symptoms associated with cachexia. Serum C-reactive protein (CRP) was agreed to be an important biomarker, but it was recognized that cachexia can be present in the absence of overt systemic inflammation [10].

In the intervening years with greater knowledge of the importance of systemic inflammatory responses in the progressive nutritional and functional decline of patients with cancer, this statement has been increasingly called into question and measurement of the magnitude of the systemic inflammatory is now integral to the definition and treatment of cancer cachexia [11–14]. This more nuanced definition reflects the evolution of criteria in the definition of malnutrition in which cancer cachexia is considered as part of disease related malnutrition with inflammation [15,16]. For example, approximately 40% of patients with operable colorectal cancer considered at medium or high nutritional risk (malnutrition universal screening tool – MUST [17]) had evidence of systemic inflammation (CRP >10 mg/L) [18].

4. Sarcopenia, sarcopenic obesity and myosteatosis

Patients may present for surgery with a range of underlying nutritional syndromes and phenotypes, such as malnutrition, sarcopenia, cachexia, obesity and myosteatosis. Furthermore, these phenotypes are associated with worsened postoperative outcome.

However, screening for such syndromes is not necessarily performed routinely in clinical practice, and there is no one screening tool that is capable of distinguishing one syndrome from another [19].

4.1. Sarcopenia

A recent study showed that the surgical population in the UK tends to be older than the general population, and that the age gap is increasing with time. Between 1999 and 2015, the percentage of people aged 75 years or more undergoing surgery increased from 14.9% to 22.9%, and this figure is expected to increase further [20]. Sarcopenia is described as ‘the loss of skeletal muscle mass and strength as a result of ageing’. There are a number of definitions for sarcopenia, which rely on the measurement of the combination of both muscle function and muscle mass. These include the European Working Group of Sarcopenia in Older Persons (EWGSOP) [21], the International Working Group on Sarcopenia (IWGS) Sarcopenia Task Force [22], the Asian Working Group for Sarcopenia and the Foundation for the National Institutes for Health (Table 1) [10,21–25].

More recently, the term “sarcopenia” has taken on a different usage. The use of diagnostic cross-sectional computed tomography (CT) images at the third lumbar vertebral level (L3) for the simultaneous perioperative analysis of body composition has become increasingly popular [26]. In this surgical context, sarcopenia has come to mean reduced muscularity, without assessment of patient’s functional status. Rather than assessing skeletal muscle mass, this CT technique analyses cross-sectional skeletal muscle area which is then indexed to patient height to give a skeletal muscle volume. This technique also provides data on the mean skeletal muscle radiodensity, quoted in Hounsfield Units (HU), which is a surrogate marker of muscle quality and an indication of the presence of myosteatosis, as well as adiposity in terms of both visceral and subcutaneous fat cross-sectional area and indices. There is a large volume of literature linking preoperative sarcopenia in a range of different pathologies, including pancreatic surgery [27], gastric cancer surgery [28], esophageal cancer [29], liver transplantation [30] and colorectal cancer [31] to worsened clinical outcomes and overall survival. The strength of this relationship is even greater when the presence of sarcopenia is combined with obesity, i.e. low muscle volume in association with elevated body adiposity. A recent meta-analysis has examined this relationship in 2297 patients with pancreatic ductal adenocarcinoma, finding both sarcopenia and sarcopenic obesity to be associated with poorer overall survival (HR 1.49, $p < 0.001$ and HR 2.01, $p < 0.001$) [32].

However, there are problems of interpretation in the literature, often due to heterogeneity in the methodology of the studies leading to variability in results. There has been a degree of

variability in the cut-offs used for the diagnosis of sarcopenia (and myosteatosis). However, there are well validated BMI and gender-specific cut-offs available in the literature for cancer patients [33]. The validated technique uses CT-based analysis at the L3 level, as this was the level that the initial validation calculations were performed in order to extrapolate to the whole body. Recently, several studies have looked at body composition analysis at the fourth thoracic vertebra as an alternative in patients who are undergoing a thoracic rather than abdominal procedure [34].

4.2. Myosteatosis

Myosteatosis is the infiltration of skeletal muscle by fat, into both intermuscular and intramuscular compartments. There are a multitude of different terms used synonymously with myosteatosis, including muscle quality, radiodensity, and muscle attenuation. There has been significant research interest in the impact of myosteatosis on surgical outcomes in a range of different cancer types, including periampullary [35], ovarian [36] and rectal cancer [37]. As with the relationship between sarcopenia and obesity, there also appears to be a combined effect with myosteatosis and obesity. In a series of 2100 patients undergoing elective surgery for colorectal cancer, three body composition subtypes were independent predictors of hospital length of stay; combined sarcopenia and myosteatosis (incidence rate ratio (IRR) 1.25), visceral obesity (IRR 1.25) and myosteatosis combined with sarcopenia and visceral obesity (IRR 1.58). The risk of readmission was associated with visceral obesity alone (OR 2.66, $p = 0.018$), visceral obesity combined with myosteatosis (OR 2.72, $p = 0.005$) and visceral obesity combined with both myosteatosis and sarcopenia (OR 2.98, $p = 0.038$). There is also emerging evidence that low skeletal muscle radiodensity is involved in the etiology of, or shares mechanisms with, other comorbidities such as myocardial infarction, diabetes and renal failure [38].

4.3. Cachexia

The third body composition syndrome of interest is cachexia, which occurs as a consequence of a range of diseases, including cancer, chronic obstructive pulmonary disease, cardiac failure, renal failure and rheumatoid arthritis. Cachexia is multifactorial in etiology [39]. For example, in patients with cancer, not only is the tumor a potential driver for nutritional depletion, but patients also tend to be older (hence, sarcopenic), live a sedentary lifestyle, and often have a poor diet, as well as have other comorbidities which may impact upon body composition. Recent evidence also suggests that some cancer patients may have a genetic predisposition to weight loss and low muscularity [40].

Table 1

Definitions of Sarcopenia (taken from the Society on Sarcopenia, Cachexia, and Wasting Disorders (SCWD) website).

Definition	Function	Muscle Mass
Sarcopenia and Frailty Research Specialist Interest Group (SIG) – cachexia-anorexia in chronic wasting disease [25]	Gait speed <0.8 m/s, OR other physical performance test	Low muscle mass (2SD)
European Working Group of Sarcopenia in Older Persons (EWGSOP) [21]	Gait speed <0.8 m/s; grip strength 40 kg males, 30 kg female	Low muscle mass (not defined)
IWGS Sarcopenia Task Force [22]	Gait speed <1.0 m/s, grip strength	Low appendicular lean mass (<7.23 kg/m ² in men, 5.67 kg/m ² in women)
Sarcopenia with limited mobility (SCWD) [10]	6-min walk <400 m, OR gait speed <1.0 m/s	Low appendicular lean mass/height ²
Asian Working Group for Sarcopenia [23]	Gait speed <0.8 m/s; grip strength 26 kg males, 18 kg females	Low appendicular lean mass/height ²
Foundation for the National Institutes of Health [24]	Gait speed <0.8 m/s; grip strength 26 kg males, 16 kg females	Appendicular lean mass/BMI

There have been a number of definitions of cachexia published previously [25,41–43]. However, the most accepted definition of cancer cachexia is “a multifactorial syndrome defined by an ongoing loss of skeletal muscle mass (with or without loss of fat mass) that cannot be fully reversed by conventional nutritional support and leads to progressive functional impairment” [10]. This international consensus provided diagnostic criteria which were either weight loss exceeding 5% or weight loss greater than 2% in individuals already showing depletion as marked by a BMI <20 kg/m² or the presence of sarcopenia.

The interaction and overlap between sarcopenia, myosteatosis and cancer cachexia are not well understood currently. In addition, the interaction between these skeletal muscle variants and patient adiposity and frailty are not clear and these should be the focus of research in the future.

5. The metabolic response to immobilization and surgical trauma

There are a number of different factors which contribute to the peri- and post-surgical trauma phenotype including immobilization, reduced oral intake, anesthesia, tissue damage, subsequent immune system activation and metabolic changes.

There are significant metabolic changes associated with a period of bedrest which are paralleled in the metabolic changes occurring after surgery [44] as immobilization is one of the key components of postoperative changes. These negative changes are also observed in clinical populations and sarcopenic or frail older adults [45] and include a decline in basal energy expenditure, reduced insulin sensitivity, anabolic resistance to protein nutrition, muscle strength and physical performance as well as increased risk of falls, health-related expenditure, morbidity and mortality. The larger impact of bed rest on the rate of loss of lean muscle leg mass and strength during bedrest in healthy older adults than their young counterparts is equivocal [46,47]. On the other hand, gain of muscle mass and function as a consequence of exercise requires significant regular training over an extended period of time, with evidence suggesting that 12 weeks of resistance exercise training is necessary for a 1.5 kg gain in muscle mass in older adults [45].

As the process of muscle loss requires a considerably shorter period of time in older adults, with just seven days of bedrest resulting in 1 kg loss of lean leg muscle mass, there should, therefore, be a particular emphasis on the preservation of muscle mass during periods of muscle disuse whilst older patients are in hospital. This loss of muscle mass occurs in both the type I (slow twitch) and type II (fast twitch) skeletal muscle fibers [48]. In terms of muscle strength, the initial loss of strength occurs rapidly during a period of immobilization, irrespective of the cause of immobilization. However, this loss of strength then plateaus after around 30 days.

Older adults tend to stay longer in hospitals and after discharge experience a more pronounced decrease in ambulatory function and reduced ability to complete activities of daily living. There are a number of strategies which have been recommended to reduce muscle wasting during bedrest in older adults, including resistance exercise [49], dietary interventions such as an increase in protein intake to exceed 1 g/kg body weight/day, administration of essential amino acid (EAA) mixtures [50,51], as well as the combination of these EAA mixtures with carbohydrate [52] or leucine, valine and isoleucine. A study [51] on the role of essential amino acids in older adults undergoing 10 days best rest found that although this normalized muscle protein synthesis, it did not have an effect upon skeletal muscle loss or function. However, when beta-hydroxy-beta-methylbutyrate (HMB) supplementation was used in a randomized placebo-controlled trial [46] in healthy volunteers

undergoing a period of 10 days bedrest, this resulted in a significant reduction in the amount of muscle loss associated with the bedrest as well as an increase in muscle mass gain during the 8 week rehabilitation phase, both in terms of total lean mass and total leg lean mass. Muscle strength also appeared to be preserved in this study.

There are many parallels to that associated with immobilization when bedrest as a consequence of surgery is considered. Preoperative fasting is associated with characteristic metabolic changes. After just a short overnight fast, the body remains able to cope with the glucose demands placed on it by the muscle, brain, kidney, bone marrow and lymph nodes by the breakdown of glycogen within the liver. However, after starvation of 24 h, the metabolic response changes to the breakdown of adipose tissue to mobilize fatty acids which are utilized by the muscle and kidney. When more prolonged periods of fasting are considered, the metabolic response becomes somewhat more complex. Muscle protein breakdown releases amino acids such as alanine and glutamine which are used in the kidney and liver to promote gluconeogenesis, with persistence of adipose tissue breakdown to provide ongoing energy stores.

Resting energy expenditure (REE) increases after surgery, with the degree determined by the magnitude of the insult, with most pronounced changes observed in those following major burns, followed by those with sepsis or peritonitis. Elective surgery is associated with a much lower increase in REE. The metabolic response to surgical trauma allows mobilization of glucose and glutamine to provide substrate for wound healing, and amino acids for acute phase protein synthesis. Intensive care unit stay is also associated with a typical pattern of skeletal muscle loss [53] which is far more rapid than that seen after a standard surgical insult.

Surgery results in an overall reduction in lean leg muscle mass [54]. However, when protein turnover is examined, there is not a large difference between the pre- and post-operative phases. When patients are fed postoperatively, this results in a significant increase in protein synthesis rates and reduction in protein breakdown when compared with patients who were fasted postoperatively [54]. Changes in skeletal muscle mass and function following surgery are most likely the consequence of inactivity combined with reduced food intake and specific metabolic changes.

6. Nutrition and surgical outcome – lessons from the ESPEN nutritionDay

In the nutritionDay dataset [55] (155 524 patients) 41% of the enrolled participants were surgical patients. The median length of stay for the cross-sectional nutritionDay data collection was 6 days for surgical and non-surgical patients [56]. Surgical patients were 6 years younger than non-surgical patients (63 vs. 69 years, $p < 0.001$). BMI was similar in surgical and non-surgical patients. BMI was <18.5 kg/m² in 7.1% of patients and was >30 kg/m² in 19%.

Weight loss within the last 3 months was slightly less frequent in surgical patients (39%) than in non-surgical patients (43%) ($p < 0.0001$) while stable weight was more frequent in surgical patients (40% vs. 33%, $p < 0.0001$). Reduced intake in the week before nutritionDay was slightly less frequent in surgical (44%) than in non-surgical (46%) patients ($p < 0.0001$). On nutritionDay the full served meal was eaten by only 35% of surgical patients vs 38% of non-surgical patients. Nothing was eaten by 20% of surgical patients and 11% of non-surgical patients mostly because they were not allowed to eat. The high proportion of surgical patients who had eaten nothing on nutritionDay is shown in Fig. 1 for preoperative, postoperative and non-surgical patients. Artificial nutrition was used in a minority of patients eating nothing. In patients not allowed to eat 30% received artificial nutrition, and in patients eating nothing despite being allowed to eat 27% received artificial




Fig. 1. Prevalence of decreased eating and association with 30-day hospital mortality in preoperative, postoperative and non-surgical patients. Each dot represents 1% prevalence within the patient group. Normal eating is shown in green and is the reference category for calculation of the univariate odds ratio for death in hospital within 30 days shown as estimate with 95% confidence intervals. (For interpretation of the references to color in this figure legend, the reader is referred to the Web version of this article.)

nutrition. Reduced eating was associated with a delay in discharge of about 1 day. Outcome at day 30 after nutritionDay was available for 83% of patients. Most patients (72.5%) were discharged home 3.8% had died in hospital. Mortality was lower in surgical patients (2%) when compared with non-surgical patients (5%).

Weight loss was associated with a slightly higher odds ratio for death in hospital within 30 days in surgical patients when compared with non-surgical patients (OR 3.2 vs 2.5). Reduced intake in the previous week was associated with a progressive increase in death within 30 days from OR 2.0 for less than normal eating, OR 3.6 for eating half and OR 6.4 for eating less than a quarter. This association was similar at all levels for non-surgical patients. Eating half the recommended amount in hospital on nutritionDay was associated with an OR 2.3 of death whereas eating nothing despite being allowed to eat was associated with an OR 9.0 (Fig. 1).

7. The patient at risk and nutritional assessment

The German hospital malnutrition study [57] found that overall 27.4% of patients were diagnosed with malnutrition according to the subjective global assessment (SGA), with a huge degree of variability between specialties. In patients who had undergone major abdominal surgery the prevalence of malnutrition was 44%, with lowest rates in those undergoing chest or general surgery (20% and 14%, respectively). A study of 26 hospital departments spread across the European Union using the nutritional risk screening (NRS-2002) tool identified that 32.6% of patients were at 'high risk' of malnutrition, with these patients developing more complications (30.6% vs 11.3% $p < 0.001$), increased mortality rates (12% vs. 1%, $p < 0.001$) and longer hospital length of stay (median 9 vs. 6 days, $p < 0.001$) when compared with patients who were 'not-at-risk'. A progressive degree of malnutrition, from none to severe, has been associated with progressive increased risk of morbidity and mortality as well as increased ICU admission and overall hospital length of stay in patients undergoing liver transplantation [58]. This relationship of increased morbidity and mortality amongst those

with malnutrition is also seen in those undergoing abdominal surgery for cancer [59].

Many of the screening tools used historically to identify those at high risk of malnutrition considered only single parameters. However, these do not facilitate the identification of patients' preoperative nutritional status, nor do they precisely identify those at high nutritional risk [60]. A validated screening tool offers a far superior method for identifying those at high risk of malnutrition correctly. Four central criteria were proposed to identify those at high nutritional risk; body mass index (BMI) and a detailed nutritional history, the presence of pathological weight loss, appetite and food intake and the severity of the underlying disease. This led to the development of a range of screening tools including the malnutrition screening tool (MST), the malnutrition universal screening tool (MUST) [17], the nutrition risk index (NRI) [61], the subjective global assessment (SGA) [62], the mini nutritional assessment short form (MNA-SF) [63] and the nutritional risk screening (NRS-2002) [64]. There is only expert consensus regarding the best screening tool available for nutritional risk assessment, which suggests that the MUST is superior in the community, NRS 2002 for inpatients and SF-MNA for those in older adult care homes. A multitude of studies have subsequently been performed to validate the predictive value for complications and mortality of preoperative NRS 2002 in patients undergoing surgery, including gastric cancer surgery [65], colorectal surgery [66] and major gastrointestinal surgery [67,68]. A meta-analysis [69] examining the use of NRS 2002 as a predictor of postoperative outcomes in abdominal surgery included a total of 11 studies. Postoperative complications were more frequent in those deemed 'at risk' than those 'not-at-risk' (OR 3.13, $p < 0.00001$). Mortality was also higher in patients 'at risk' (OR 3.61, $p = 0.009$) and these patients had a significantly longer hospital LOS (mean difference 3.99 days, $p = 0.01$) [69].

More recent guidelines [1] have explored criteria for the diagnosis of severe nutritional risk, and these have included weight loss exceeding 10–15% within the preceding 6 months, BMI less than 18.5 kg/m², NRS 2002 > 5 or SGA grade C or a preoperative serum

albumin concentration less than 30 g/L in the absence of hepatic or renal dysfunction. If one of these criteria is present, targeted nutritional therapy should be instigated immediately. If the screening tools discussed previously identify a patient at risk, a more formal and extensive nutritional assessment should be performed by an appropriately trained professional. This assessment should include nutritional assessment using a plate chart or 24-h dietary recall, estimation of patients subcutaneous and visceral adiposity and skeletal muscle mass, other anthropometrics measures such as upper arm circumference and skin-fold thickness; hand-grip strength as a test of muscle function; and Barthel index or 6-min walking test as a measure of body function [70].

8. Preoperative nutritional and metabolic preparation of the surgical patient

Preoperative conditioning is defined as the process of training to become physically fit by a regimen of exercise, diet and rest and is, therefore, regarded as a multimodal intervention. Perioperative oral nutrition is considered one of the major preoperative components of Enhanced Recovery After Surgery (ERAS) pathways [71]. ERAS is believed to help by 'exploiting the critical perioperative period to improve long-term cancer outcomes' [72], and optimization of nutrition is one area which can be exploited successfully.

The concept of preoperative conditioning is not a new one. In 1992 the concept of a 'decision box' [73] which helps to identify the right patients who will benefit most from a nutritional intervention, was devised. Given the high prevalence of malnutrition discussed in the previous section and the known risk factors, which are highly prevalent amongst those undergoing surgery, this should be aggressively targeted. The metabolic risk is exacerbated in patients with malignancy [74] due to release of TNF- α , IL-6 and IL-1 in addition to anorexia caused by central nervous system signaling which results in muscle wasting, changes in liver metabolism as well as consumption and depletion of fat stores. Exercise is one modality which can help modulate these metabolic consequences of tumor, by promoting IGF-1, mTOR, and Akt which results in increased protein synthesis; IL-10, sTNF-r1 and sTNF-r2 which reduces systemic inflammation; GLUT-4 which reduces insulin resistance; and superoxide dismutase and glutathione which results in a reduction in the formation of reactive oxygen species [75].

The aims of preoperative conditioning are to restore the energy deficit, improve functional performance, avoid weight loss and preserve the gut microbiome. To obtain such effect, a normocaloric diet is sufficient with a protein intake of 1.2 g/kg [76]. The intervention should include dietary counselling, fortified diets, oral nutritional supplementation (ONS), and parenteral support, where indicated. The enteral route is always preferred wherever feasible and even when patients are consuming a normal diet this is frequently insufficient to obtain their energy requirement, so it is recommended that patients receive oral nutritional supplements (ONS) in the preoperative period, irrespective of their nutritional status [1]. There is good evidence to support ONS in the perioperative period, with a meta-analysis of 9 studies [77] finding this to be associated with a 35% reduction in total complications ($p < 0.001$) and this translated to a cost saving and was also found to be cost effective. In those patients who are identified as high-risk undergoing major abdominal surgery and those who are malnourished with a diagnosis of cancer, ONS should be considered obligatory [1]. In terms of parenteral nutrition (PN), this should only be considered in those with malnutrition or severe nutritional risk where emergency requirements cannot be met by enteral nutrition interventions alone [1]. Where this approach is absolutely necessary, PN should be provided for 7–14 days preoperatively to maximize

benefit, based upon evidence that this time frame is necessary to reduce the Clavien-Dindo grade 3b or higher surgical site infection-based complications [78].

The use of carbohydrate loading as metabolic conditioning is supported by some basic science and clinical studies [79,80]. A recent large prospective randomized clinical trial (RCT) has shown significant benefits regarding the reduction of postoperative insulin resistance and hyperglycemia without impact on the complication rate [81]. So far, the evidence for a decrease of postoperative morbidity is not yet clear.

Prehabilitation has gained popularity in recent times, with increasing evidence to support a multimodal prehabilitation program in a range of surgical specialties. A study combining a 6-week preoperative bundle of physical exercise and endurance training, nutrition interventions and psychological support to improve anxiety when compared with postoperative rehabilitation alone [82] in a cohort of patients undergoing elective colorectal surgery found that this optimizes the patient's functional capacity throughout the perioperative period. In those patients who are due to undergo preoperative neoadjuvant therapy, the period after cessation of therapy but prior to surgery is typically 4–6 weeks and this time should be exploited to optimize patient fitness. A meta-analysis of multimodal prehabilitation [83] in elective colorectal surgery found that this was associated with a significant reduction in hospital length of stay of 2 days and was linked to a faster time to return to presurgical functional capacity. When pooled data from RCTs regarding trimodal prehabilitation was analyzed [84], this found that the postoperative loss of lean body mass was attenuated in patients undergoing prehabilitation versus rehabilitation alone. There is also evidence that a multimodal intervention is associated with improved perioperative physiological parameters, functional outcomes and quality of life measures, but no impact on postoperative complications in those undergoing liver resection [85] as well as for a beneficial effect in muscle strength in sarcopenic older adult patients undergoing gastric cancer resection [86]. In high-risk patients undergoing elective major abdominal surgery, a RCT found that prehabilitation in the form of a motivational interview, high-intensity endurance training and promotion of physical activity was associated with a significant reduction in the incidence of postoperative complications (31% vs. 62%, $p = 0.001$) [87].

9. Perioperative glycemic control

Hospital guidelines surrounding perioperative glycemic control are based, in 90% of cases, on the guidance published by Diabetes UK in 2011 [88]. This provides a standard of care, which should be met commencing at the point of referral from primary care, through the perioperative phase and to discharge from hospital. At the first stage when the patient is referred from primary care, the minimum information that should be provided should include the duration and type of diabetes, the place of usual diabetes care (primary or secondary), other comorbidities, and treatment (both for the diabetes and other comorbidities). Information should also be provided on details of any diabetes-associated complications such as renal or cardiac disease, and finally any relevant measures from within the last 3 months, including body mass index (BMI), blood pressure, HbA1c and estimated glomerular filtration rate (eGFR). However, the compliance to this standard was low [89].

There is evidence supporting an association between the presence of diabetes and significantly elevated risk of 30-day mortality in patients undergoing elective non-cardiac surgery [90]. Those patients with diabetes (20.2%) with preoperative hyperglycemia (7.9%) were twice as likely to die as those with a normal preoperative glucose concentration. However, if the patient did not have preoperatively diagnosed diabetes but had preoperative




Fig. 2. Suggested algorithm for perioperative fluid therapy.

hyperglycemia, they were 13 times more likely to die within 30 days of surgery when compared with a patient with normal preoperative glucose concentration. When postoperative hyperglycemia was considered, if patients were not diagnosed with diabetes but had perioperative or postoperative hyperglycemia, they were 45 times more likely to die than those with a normal glucose concentration. There is also an association between hyperglycemia in those who were previously normoglycemic and composite adverse events [91], as well as reoperative interventions, anastomotic failures, myocardial infarction and composite infections [92]. However, knowing that the patient was diabetic in the presence of hyperglycemia attenuated these worse clinical outcomes by almost half. There is consistent evidence that the highest risk group with regards to perioperative glucose control are those who are not diagnosed with diabetes but who develop postoperative hyperglycemia.

Clinical outcomes in those with poorly controlled diabetes are significantly worse than those with well-controlled diabetes, with a stepwise increase in the risk of infectious complications and mortality relating to infection according to increasing HbA1c (RR 0.98, if HbA1c <6% vs. RR 2.01, if HbA1c \geq 11%) [93]. Patients with highest preoperative HbA1c levels tend to have their blood glucose concentrations checked earlier, have higher postoperative glucose concentrations and are significantly more likely to be commenced on insulin postoperatively, than those with a lower preoperative HbA1c, possibly due to an elevated level of vigilance [94].

The current National Institute for Health and Care Excellence (NICE) clinical guideline 45 on the use of routine preoperative tests prior to elective surgery suggests that HbA1c should only be routinely tested in those patients with a formal diagnosis of diabetes [95]. However, this is a controversial policy as it fails to identify those patients with non-diabetic hyperglycemia [96] and, therefore, misses the opportunity to intervene preoperatively and modify the elevated perioperative surgical risk that this is associated with.

10. Perioperative fluids and outcome

There is a close relationship between nutrition and fluid and electrolyte balance, with the intake of food by natural or artificial means being inseparable from that of fluid and electrolytes [97]. The metabolic response to surgery is associated with salt and water retention and an increase in the excretion of potassium, as a result of which patients are susceptible to retention of salt and water, and

consequent fluid overload in the perioperative period [98–103]. There is a relatively narrow margin of safety in perioperative fluid therapy and either too much or too little can have a negative effect on physiological processes and clinical outcome. The goal of perioperative intravenous fluid therapy should be to maintain tissue perfusion and cellular oxygen delivery, while at the same time keeping the patient in as near zero fluid and electrolyte balance as possible (Fig. 2).

10.1. Preoperative period

Patients should reach the anesthesia room in a state as close to euvoolemia as possible with any preoperative fluid and electrolyte imbalance having been corrected. Current anesthetic recommendations that allow patients to eat for up to 6 h and drink clear fluids up to 2 h prior to the induction of anesthesia help to prevent preoperative fluid depletion without increasing aspiration-related complications. Some patients may need intravenous fluids to restore euvoolemia prior to surgery.

10.2. Intraoperative period

Most patients require crystalloids at a rate of 1–4 ml/kg/h to maintain homeostasis [104]. However, some patients develop intravascular volume deficits which require correction by administration of goal-directed boluses of intravenous solutions. Goal directed fluid therapy (GDFT) is aimed at maintaining intravascular normovolemia guided by changes in stroke volume as measured by a minimally invasive cardiac output monitor to optimize the position of each patient on his/her individual Frank–Starling curve [105,106]. In addition to the background crystalloid infusion, fluid boluses (200–250 ml) should be given to treat any objective evidence of hypovolemia (>10% fall in stroke volume) in order to optimize intravascular volume and cardiac output [107]. A recent meta-analysis that included 23 studies with 2099 patients has shown that GDFT was associated with a significant reduction in morbidity, hospital length of stay, intensive care length of stay, and time to passage of feces [108]. However, when patients were managed within ERAS pathways, with optimal perioperative care and avoidance of postoperative fluid overload, the only significant reductions were in length of intensive care stay and time to passage of feces. It has also been shown that GDFT does not impact on outcome when compared with conventional intraoperative fluid therapy in patients undergoing elective colorectal surgery [109].

Hence, within ERAS programmes, it may not be necessary to offer all patients GDFT, which should be reserved for high-risk patients or for patients undergoing high-risk procedures [104].

10.3. Postoperative period

For most patients undergoing elective surgery, intravenous fluid therapy is usually unnecessary beyond the day of operation, except for those undergoing upper gastrointestinal and pancreatic procedures. With these exceptions, patients should be encouraged to drink as soon as they are awake and free of nausea after the operation. An oral diet can usually be started on the morning after surgery [110,111]. When adequate oral fluid intake is tolerated, intravenous fluid administration should be discontinued and be restarted only if required to maintain fluid and electrolyte balance. If intravenous fluids are required, then in the absence of ongoing losses, only maintenance fluids should be given at a rate of 25–30 ml/kg/day with no more than 70–100 mmol sodium/day, along with potassium supplements (up to 1 mmol/kg/day) [112]. As long as this volume is not exceeded, hyponatremia is very unlikely to occur despite the provision of hypotonic solutions [113,114]. Any ongoing losses (e.g. vomiting or high stoma losses) should be replaced on a like-for-like basis, in addition to maintenance requirements. After ensuring the patient is normovolemic, hypotensive patients receiving epidural analgesia should be treated with vasopressors rather than indiscriminate fluid boluses [115,116]. Fluid deficit or overload of as little as 2.5 L [117] can cause adverse effects in the form of increased postoperative complications, prolonged hospital stay and higher costs due to increased utilisation of resources [118–120].

An excess of 0.9% saline causes hyperosmolar states, hyperchloremic acidosis [121–126], and decreased renal blood flow and glomerular filtration rate, which in turn exacerbates sodium retention. Edema impairs pulmonary gas exchange and tissue oxygenation leading to an increase in tissue pressure in organs such as the kidney which are surrounded by a non-expandable capsule. Microvascular perfusion is compromised, arterio-venous shunting increases and lymphatic drainage is reduced, leading to further edema. Hyperchloremic acidosis, as a result of saline infusions has been shown to reduce gastric blood flow and decrease gastric intramucosal pH in older adult surgical patients, and both respiratory and metabolic acidosis have been associated with impaired gastric motility. Fluid overload also causes splanchnic edema resulting in increased abdominal pressure, ascites and even the abdominal compartment syndrome, which may lead to decreased mesenteric blood flow and ileus, with delayed recovery of gastrointestinal function, an increase in gut permeability, intestinal failure and even anastomotic dehiscence [127].

Fluid restriction resulting in fluid deficit can be as detrimental as fluid excess by causing decreased venous return and cardiac output, diminished tissue perfusion and oxygen delivery and increased blood viscosity. It can also lead to an increase in the viscosity of pulmonary mucus and result in mucous plug formation and atelectasis [128]. Induction of anesthesia in patients with a fluid deficit further reduces the effective circulatory volume by decreasing sympathetic tone. Inadequate fluid resuscitation and decreased tissue perfusion can lead to gastrointestinal mucosal acidosis and poorer outcome.

A meta-analysis of patients undergoing major abdominal surgery has shown that patients managed in a state of near-zero fluid and electrolyte balance had a 59% reduction in risk of developing complications when compared with patients managed in a state of fluid imbalance (deficit or excess). There was also a 3.4-day reduction in hospital stay in the near-zero fluid balance group [120].

11. Inflammation and surgical outcome

The “trauma of surgery” leads to release of stress hormones and inflammatory mediators. This so-called metabolic stress is akin to the “Systemic Inflammatory Response Syndrome” (SIRS) that follows any injury or infection and is mediated by cytokines. This syndrome induces catabolism of stores of glycogen, fat and protein leading to release of glucose, free fatty acids and amino acids into the circulation – to support the process of tissue healing. It is, therefore, important to have sufficient protein reserves, preoperatively. This is because current thinking is that, whilst postoperative nutritional therapy may provide the energy for optimal healing and recovery, in the immediate postoperative phase it may only minimally counteract muscle catabolism, or not at all [1]. The consequences of insufficient protein reserves in the postoperative patient include: decreased wound healing, impaired immune response, defective gut-mucosal barrier and decreased mobility and respiratory effort. All of these would lead to an overall poorer postoperative course [129].

11.1. Systemic inflammatory response (SIR)

As described in the American Critical Care Medicine consensus [130], SIRS is described by any two of the following: a temperature $>38\text{ }^{\circ}\text{C}$ (100.4°) or $<36\text{ }^{\circ}\text{C}$ ($96.8\text{ }^{\circ}\text{F}$); heart rate >90 beats/min; respiratory rate >20 breaths/min or $\text{PaCO}_2 <32$ mmHg; white blood cells $>12 \times 10^9$ cells/l or $<4 \times 10^9$ cells/l or $>10\%$ immature (band) forms [130] as well as the absence of a source of an infective focus [130]. In addition to this definition there many pathophysiological changes that occur as part of the systemic inflammatory response (Table 2) [131].

11.2. The importance of C-reactive protein (CRP)

The prototypical marker of the systemic inflammatory response is CRP. A systematic review that explored routine clinical markers and their association to the magnitude of systemic inflammatory response (SIR) after surgery found that even though cortisol, IL-6, WCC, and CRP all peak after all types of elective operations (minor and major, laparoscopic and open), only IL-6 and CRP were consistently associated with the magnitude of the operative injury [132]. CRP is routinely measured in clinical laboratories world-wide and used extensively in clinical practice and, therefore, may be useful in the monitoring and modulation of the SIR after elective operation. A systematic review and meta-analysis that included 22 studies, of which 16 studies were eligible for meta-analysis, found that the pooled negative predictive value (NPV) of CRP improved each day after surgery up to

Table 2
Pathophysiological changes of the systemic inflammatory response.

Neuroendocrine changes
Fever, somnolence, fatigue and anorexia
Increased adrenal secretion of cortisol, adrenaline and glucagon
Hematopoietic changes
Anemia
Leucocytosis
Thrombocytosis
Metabolic changes
Loss of muscle and negative nitrogen balance
Increased lipolysis
Trace metal sequestration
Diuresis
Hepatic changes
Increased blood flow
Increased acute phase protein production

Table 3
Systemic inflammation and its effects on the surgical patient.

Protein catabolism after surgery leads to depletion of lean body mass.
The magnitude of the postoperative systemic inflammatory response corresponds to the amount of surgical trauma.
The higher the response the poorer the surgical outcome.
C-reactive protein is useful in quantifying the magnitude of the postoperative systemic response.
Moderating the postoperative systemic inflammatory response (e.g. by using a laparoscopic approach) appears to improve surgical outcome.

90% at postoperative day (POD) 3 for a pooled CRP cutoff of 159 mg/L [133], and concluded that infectious complications after major abdominal surgery are very unlikely in patients with a CRP below 159 mg/L on POD 3 [134]. Another systematic review and pooled-analysis evaluating the predictive value of CRP for major complications after major abdominal surgery calculated a prediction model based on major complications as a function of CRP levels on the third postoperative day [135]. Based on the model a two cut-off system was suggested consisting of a safe discharge criterion with CRP levels below 75 mg/L and above 215 mg/L serving as a predictor of complications [135].

This work highlights the clinical utility of CRP to identify the magnitude of the effect of surgery on postoperative protein catabolism and clinical outcomes. Also, CRP provides an indicator on which to judge the effect of interventions to mitigate the effects of the SIR in the postoperative period. In this context there is good evidence to support the use of laparoscopic surgery [136] and preoperative steroids [137]. Also, there is some evidence that supports the use of preoperative oral antibiotics in combination with mechanical bowel preparation [138,139].

The importance of systemic inflammation and its effects on the surgical patient are summarized in Table 3.

12. The impact of enhanced recovery after surgery

Enhanced Recovery After Surgery (ERAS) is a relatively new pathway of care for the surgical patient [140]. It is a multi-modal, multi-disciplinary and evidence-based approach to perioperative care, where teams of professionals work together to achieve best practice at all times, but also are ready and able to adapt and adopt new improvements.

The first evidence-based guidance for the entire perioperative care of a patient undergoing major surgery was published in 2005 [71]. The literature showed clear evidence of benefit for avoiding bowel preparation, wound drains, nasogastric tubes, removing urinary catheters, stopping intravenous fluids early and allowing early feeding. Modern fasting guidelines allowing drinking of clear fluids two hours before surgery, and avoiding long acting pre-medication. Long-acting anesthetic agents and opioids for pain management should be avoided (Fig. 3). All these treatments had good evidence for their use but were rarely practiced at that time. The evidence is constantly being updated and recommendations may change as the evidence base increases. This is exemplified by that fact that although mechanical bowel preparation on its own is of no benefit [141], the combination of oral antibiotics and mechanical bowel preparation may reduce surgical site infections and anastomotic leaks [139].

However, it was found that a protocol on its own was not enough. The care around the patients and the hospital management infrastructure needed to be organized differently [142]. First of all, there is a need to audit what is actually being done with regard to all the recommended ERAS care elements. The patient is passing through several units and different departments during the care process. In each one of these, many professionals are managing their specific focuses for the time they have the patient to care for. Once done, they pass the patient over to the next care giver. The complexity of the organization is such that no one has any overview or full control of the entire care pathway. This was a primary need that was addressed by the ERAS group by instituting audit for each and every patient.

Since the patient is treated by many different professionals and they work in different parts of the hospital, it was necessary to form teams that covered all stations and all professions. This led to the birth of the ERAS Team. This team is led by doctors from surgery and anesthesia who take the medical responsibility for the care that is delivered and administered and run by nurses, led by an ERAS coordinating nurse.

A major breakthrough for ERAS came in 2010 when it was reported in meta-analysis that ERAS reduced complications [143]. Now the data suggest 50% reductions in complications after colorectal surgery. This sparked a lot of interest and soon ERAS principles were employed for most major operations in RCTs and case series, all showing similar outcomes with faster and better recovery




Fig. 3. Elements of Enhanced Recovery After Surgery pathways in the pre-, intra- and post-operative periods.

[144]. This also held true for the most vulnerable patient groups such as the frail and older adults [145]. ERAS also reduces the impact of risk factors including diabetes [146], undernutrition [147] and facilitates optimal metabolic and nutritional care [148].

When ERAS is combined with minimally invasive surgery poor compliance to the protocol may overshadow the risks associated with co-morbidities [149]. The main mechanisms behind these improvements are likely to be associated with the marked reduction in stress reactions to the surgery, since many of the elements of ERAS have this effect [150]. In colorectal surgery, better compliance with the protocol results in shorter stay, fewer readmissions, fewer complications [151,152] and is associated with improved 5-year survival [153].

The variation in care delivery and outcomes is huge worldwide [154], within continents, in countries [155] and between different practitioners [156,157]. Much of this variation is due to the slow adoption of modern care and the practice of old and outdated care principles. The reasons for this are many, but it is interesting to find that the implementation program run by the ERAS Society has proven to work in all major continents and in different socio-economic environments. With the marked reduction in complications and the opening up of resources with faster recovery and shorter stay the economics of ERAS is positive regardless of financing of the health care system [158].

In summary, the evidence-based multi-modal and multi-professional approach to perioperative care – ERAS – has been shown to markedly improve surgical outcomes and reduce the costs of care.

13. Recovery in the community

Following a successful perioperative hospital stay, setting of expectations and thorough preparation are key to a successful discharge from hospital including pain management, nutrition, the use of laxatives for return of bowel function, appropriate exercises to help regain normal function, and having a contact point for any questions. Information should also be provided about symptoms to be wary of, which may indicate the presence of a complication, and what to expect in terms of follow-up. There is good evidence that nursing telephone follow-up following discharge is positive in terms of providing support and reassurance for patients [159], as well as reducing hospital readmission rates and improving patient satisfaction. The process of expectation setting commences with preoperative counselling [160] where the patient is provided information regarding what to expect on a daily basis after surgery, identifying the resources available to the patient to facilitate smooth recovery, and what the patient can do to optimize their outcome. This information giving is frequently backed up with comprehensive guides and booklets to help them better understand ERAS programs. In terms of post-discharge from hospital, support from the district nurse or home helper is invaluable in providing information regarding adequate nutrition, continued rehabilitation and exercise.

14. Postoperative nutrition

The instigation of postoperative nutrition should be a part of routine care rather than an afterthought. In addition, ensuring establishment of early oral nutrition is a fundamental tenet of ERAS [1].

The mode of nutritional delivery in the early postoperative period has been a subject of much debate, especially in procedures involving the formation of bowel anastomosis. However, several studies and systematic reviews with meta-analysis have concluded that the oral and/or enteral route is the preferred mode of nutrition

for surgical patients. A review of five feeding routes following pancreaticoduodenectomy showed that nutritional delivery via the oral route was associated with the least complications [161]. A more recent meta-analysis using only RCTs showed enteral to be superior to parenteral nutrition following pancreaticoduodenectomy [162].

Avoidance of oral intake, which was felt to reduce the risk of complications, especially after gastrointestinal surgery involving anastomoses has not been demonstrated to be beneficial in the setting of RCTs. However, this avoidance of nutritional intake carries the very real risk of postoperative underfeeding of an already at risk patient group. This could further exacerbate malnutrition and influence postoperative complication rates.

There is a distinct requirement of the understanding of this metabolic response and how to optimize or support the post-operative patient with the appropriate nutritional therapy especially in instances when the patient is malnourished. The long term caloric and protein deficits in the postoperative patient results in poorer postoperative outcomes.

14.1. Early postoperative nutrition

Early nutrition has been shown in abdominal and pelvic surgery to stimulate peristalsis and gastrointestinal function, reduces the risk of postoperative ileus and shorten overall hospitalization period. It was observed that patients who had earlier enteral feeding had fewer complications after colorectal surgery (4.5%) vs late enteral nutrition (19.4%) [163]. A Cochrane review on early enteral nutrition also showed no difference in risk of postoperative complications in patients fed early (within 24 h) and those fed late. Importantly they showed that patients who were fed early had a reduction in mortality RR (0.41, 95% CI 0.18 to 0.93) [164]. An updated review on the same premise found reduction in length of hospital stay but was inconclusive on postoperative outcomes and quality of life [165].

14.2. Routes of feeding

The current ESPEN guidelines state that 'Oral nutritional intake shall be continued after surgery without interruption and oral intake, including clear liquids, shall be initiated within hours after surgery in most patients' [1]. Perioperative nutritional support therapy is indicated in patients with malnutrition and those at nutritional risk. Perioperative nutritional therapy should also be initiated, if it is anticipated that the patient will be unable to eat for more than five days perioperatively. It is also indicated in patients expected to have low oral intake and who cannot maintain above 50% of recommended intake for more than seven days. In these situations, it is recommended to initiate nutritional support therapy without delay.

This is further supported by the systematic reviews and meta-analyses on several gastrointestinal surgical procedures that have shown no increased benefit of food avoidance and indeed better outcomes in patients who received oral nutrition and those who were fed enterally [161,164–166]. In all of these instances they found that early enteral and oral nutrition was not associated with an increase in clinically relevant complications, but rather a shorter length of hospital stay [161,162,165,166]. Only in cases where the energy and nutrient requirements cannot be met by oral and enteral intake alone (<50% of caloric requirement) for more than seven days, a combination of enteral and parenteral nutrition is recommended [1].

15. Postoperative exercise intervention

Exercise stimulates muscle capillarization, protein synthesis, insulin sensitivity and mitochondrial function and proliferation and, therefore, is a good strategy to maximize postoperative recovery. However, robust voluntary exercise intervention post-operatively at a time when metabolic dysregulation and fatigue are at their greatest is unlikely to be practicable, and fatigue may persist for many weeks after surgery [167]. Furthermore, muscle wasting and deconditioning will be exacerbated by prolonged periods of bed rest [44]. In this situation, non-voluntary, transcutaneous, electrically evoked muscle contraction may be an effective strategy for the maintaining or improving muscle mass and function after surgery until voluntary exercise, which is likely to be most effective, is practicable [168]. Given muscle mass restoration following wasting is known to be slower and of less magnitude in older people [169], resistance exercise intervention in older people will need to be supervised and intensive to be successful. Patient muscle mass restoration may be augmented if exercise intervention is combined with protein nutrition, although this is controversial providing the volunteer is in protein balance [170].

16. The role of novel nutrients and substrates

In the last decades, standard enteral and parenteral formulae have been supplemented with specific nutrients and substrates with the goal of improving several metabolic pathways, which are deranged by surgical injury. The peculiar and unique mechanisms of action of some substrates, established first in experimental settings, encouraged the induction of clinical trials.

16.1. Glutamine

Glutamine is involved in a variety of biological processes, such as anabolic functions, acid-base regulation in the kidney, and ammonia metabolism [171]. Depletion in glutamine storage during stressful events [172] has been reported, and exogenous supplementation is associated with improved protein synthesis, preservation of gut barrier, enhancement of wound healing, reduction of oxidative stress, negative nitrogen balance, improvement of glucose metabolism, and modulation of the immune system [173].

Until 2007, several randomized, but underpowered, RCTs had been published and when the results were pooled in a meta-analysis [174], the effect of parenteral or enteral glutamine supplementation resulted in a significant reduction of surgical morbidity and duration of hospitalization. In 2009, the largest RCT ($n = 428$) on the impact of the parenteral glutamine supplementation (0.4 g/kg/day) in major abdominal operations for cancer, rejected the hypothesis of a protective effect on any type of surgery-related morbidity and on the length of hospital stay [175]. More recently a multicenter double-blind RCT was reported including 150 surgical ICU patients without renal or hepatic failure, or shock. All received isonitrogenous isocaloric parenteral nutrition (1.5 g/kg/day). In the intervention group, glutamine was administered in the standard dosage of 0.5 g/kg/day. No significant differences were seen with the primary endpoints of hospital mortality and infection rate (mortality glutamine vs. standard 14.7% vs. 17.3%, bloodstream infection rate 9.6 vs. 8.4 per 1000 hospital days) [176].

A recent meta-analysis [177], included 19 RCTs with 1243 patients scheduled for elective major abdominal surgery. Glutamine supplementation did not affect overall morbidity (RR = 0.84; $p = 0.473$) and infectious morbidity (RR = 0.64; $p = 0.087$). Patients treated with glutamine had a significant reduction in length of hospital stay.

16.2. Omega-3 fatty acids

Fatty acids are potent modulators of the immune and inflammatory responses. They are incorporated into the cell membrane influencing the function and structure. By penetrating into the cell cytoplasm, fatty acids affect the synthesis of eicosanoids, cytokines and several other key mediators. Furthermore, they impact on gene expression and cell signaling. In addition, the cell-mediated immune responses are deeply affected by different type of fatty acids. Specifically, omega-3 fatty, as opposite to omega-6 fatty acids, stimulate the synthesis of less proinflammatory leukotrienes, prostaglandins, and thromboxanes [178].

Despite the strong molecular background, robust clinical studies on the effect of parenteral formulae containing omega-3 fatty acid-based lipid emulsion are limited. The largest RCT on this topic showed no significant difference between treatment and control arms in postoperative complication rates with an associated and unexplained 5-day reduction in LOS in the group receiving omega-3 fatty acids [179]. A recent systematic review and meta-analysis collected 49 RCTs addressing the impact of omega-3 fatty acids on surgical outcomes [180], but only 24 studies, with a total of 2154 patients, reported the rate of post-operative infections. Regardless of the commercial formulation used, the risk ratio was in favor of the group receiving omega-3 fatty acids (RR = 0.60; 95%CI [0.490, 0.72]). As properly emphasized by the authors, the major constraint of this meta-analysis [180], as well as others [181], was the inclusion of underpowered and non-significant trials. This limitation could have produced overstating results.

16.3. Enteral feeds containing multiple substrates

Most of the evidence suggesting that specific nutrients may modulate the clinical course of patients undergoing major operations has been produced by testing enteral or oral formulae enriched with arginine, omega-3 fatty acids and ribonucleotides [182,183].

The evidence has been extensively argued and reported in the 2017 ESPEN guideline on clinical nutrition in surgery [1]. The author recommendations were as follows: “peri- or at least post-operative administration of specific formulae enriched with immunonutrients should be given in malnourished patients undergoing major cancer surgery. There is currently no clear evidence for the use of these formulae enriched with immunonutrients versus standard oral nutritional supplements exclusively in the preoperative period”. These statements were based after the authors’ systematic search for studies and reviews published between 2010 and 2015. However, a recent focused meta-analysis on preoperative immune modulating nutrition in gastrointestinal cancer only, has demonstrated a significant reduction in infectious complications and tendency to a shorter length of stay [182].

It should be highlighted that the vast majority of the published RCTs on immunonutrition in surgical patients were conducted outside the implementation of ERAS protocols. The beneficial effect of the administration of immunonutrients, in addition to ERAS pathways has been addressed in recent multicenter Spanish RCT [184]. They studied this association in well-nourished patients undergoing colorectal resection for cancer. The findings demonstrated a decrease in the total number of complications observed in the immunonutrition treated group compared with the control group, primarily due to a reduction in infectious complications (23.8% vs. 10.7%, $P = 0.0007$). These findings look promising but necessitate future confirmations.

17. Pre-, pro- and synbiotics in the surgical patient

Probiotics, as defined by the World Health Organization, are live microorganisms which, when administered in adequate amounts, confer a health benefit on the host. They survive transit through the gastrointestinal tract with the majority of their activity being in the colon [185]. Prebiotics are carbohydrate compounds, primarily oligosaccharides which induce growth and/or activity of selective bacterial genera in the colon [186]. Combinations of prebiotics and probiotics in a single preparation are referred to as synbiotics [185]. Current literature suggests that multispecies preparations are more effective due to better survival of the gastro-duodenal passage or greater ability to find a biological niche. However, to date, the most appropriate species of probiotic has not been described in the currently available literature.

Probiotics have been used in the treatment of several abdominal complaints. They have been shown to be useful in the treatment of gastrointestinal infections, for oral rehydration therapy in treating acute infectious diarrhea in children [187–190], traveller's diarrhea [191] and antibiotic-associated diarrhea in both children [192–194] and adults [195–198]. Recent ESPEN guidelines stated that use of a specific probiotic multi strain mixture may be beneficial for primary and secondary prevention of pouchitis in patients with ulcerative colitis who have undergone colectomy and ileo-anal pouch anastomosis. There are some data to confirm the use of the same multi strain probiotic mixture for the treatment of pouchitis after antibiotic treatment failure as well as for the treatment of mild to moderate ulcerative colitis [199]. The suggested mechanisms of action include both direct antimicrobial effects as well as indirectly or competitively excluding potentially pathogenic bacteria [200]. They achieve this by producing bacteriocins which inhibit pathogenic epithelial adherence and production of virulence factor, and prevent bacterial translocation via tight junctions [200,201]. They also alter the gut microenvironment by altering the mucosal pH, which further inhibits pathogenic bacteria. Additionally, others have shown that probiotic bacteria can hamper the inflammatory response by promoting anti-inflammatory cytokine production [200,202]. Whilst these nutritional adjuncts are emerging as potential treatments that could help reduce the incidence of post-operative infection, the success or failure of one strain cannot be extrapolated to other strains.

To the post-surgical patient, the stress of the operative procedure can lead to a proinflammatory stimulus that increases gut permeability. Increased gut permeability together with dysbiosis may lead to increased bacterial translocation across the gut barrier into the circulation. Bacterial translocation is an important pathogenic factor for the increased risk of infections. To this end the introduction of probiotics or synbiotics would be expected to maintain gut barrier function by restoring intestinal permeability ameliorating the intestinal inflammatory response and the release of cytokines, and maintaining the homeostasis of the normal gut microbiota.

A number of RCTs have examined the value of prebiotics and probiotics in reducing postoperative complications in particular post-operative infective complications. The interest in synbiotics, is based on emerging evidence that the proliferation of probiotic bacteria can be enhanced by the co-administration of prebiotics [203]. Indeed a more recent meta-analysis has shown that whilst infectious complications were reduced after elective abdominal surgery, the effect was better still in those patients who received synbiotics [204].

Contrastingly, some studies have yielded mixed results that probably are due to the variations of probiotics used,

methodological quality and study endpoints. Additionally, others have described adverse events surrounding the use of probiotics. It is, however, noteworthy that serious adverse effects of probiotics are uncommon in those who are well. In patients with severe pancreatitis, administration of probiotics was associated with an increased frequency of bowel ischemia – the mechanism of this is still unexplained [205–207]. However, this effect of probiotics has not been identified in any other study. In the most recent meta-analysis [204], no serious adverse events were noted. The authors concluded that probiotics and synbiotics are safe in elective gastrointestinal surgery and is associated with a significant reduction in infectious complications.

18. Patient and caregiver partnership

The period surrounding a major surgical procedure is highly taxing on patients and their caregivers. Perioperative nutrition is recognized as a substantial issue, with significant weight loss being not uncommon. Malnutrition in this setting is multifactorial, including issues with poor appetite, unappealing hospital nutrition, postoperative pain and a reduced level of consciousness. Support from family is frequently key to optimizing perioperative nutritional intake and modification of previous eating habits including consuming high calorie foods on a little but often basis. Oral nutritional supplementation is often met with variable patient acceptability and hence, compliance is often not optimal. The effects of major surgery and indeed the complications, have wide reaching effects on not just the patient but also their families and caregivers, rendering them a bystander in the care of their loved ones. The importance of communication cannot be over-emphasized in this setting, and a strong partnership between the surgeon and patient, family and caregivers is needed to overcome complicated postoperative courses.

19. Future directions for research and policy

The evidence contains many strengths, and these are reflected in high-quality guidelines surrounding perioperative nutrition [1]. However, there are still many areas of nutrition in this setting which have not yet been fully explored. An area of research development surrounds the global obesity epidemic and its link to the metabolic syndrome, with more attention being directed towards a multidisciplinary approach to the management of obesity and its related diseases [208], tying together concepts such as bariatric and orthopedic surgery, geriatrics, endocrinology, psychology and psychiatry, as well as nephrology and dialysis. An area of research which is going to become increasingly relevant is the shift in population related to the ageing epidemic which is currently underway. With increasing frailty comes weight loss, progressive skeletal muscle weakness, exhaustion and inactivity, all of which increase the prevalence of disability, loss of independence and worsened clinical outcomes.

Not only are there challenges in developing an evidence base for interventions, but also in the implementation of this evidence once established. One area in which implementation lags behind the evidence base for its practice is ERAS protocols in surgery, with a multicenter qualitative study finding the main barriers to implementation being time restraints, a reluctance to change and the logistics of setup [209]. Another topic is that of fasting guidelines in enterally fed in critical care patients. Again, this identified issues surrounding mistrust of the guideline, resistance to a change in clinical practice, as well as perceived increased clinical complexity which all acted as barriers to implementation. There are some key

Table 4

Take home messages.

- History is continuity – those who don't learn from the lessons of history are condemned to repeat it.
- Preoperative muscle mass is critical to postoperative outcome.
- Sarcopenic obesity is an independent predictor of postoperative complications, especially when the host genotype is associated with weight loss and a low skeletal muscle index.
- Surgical patients who don't eat when eating is allowed have an increased length of stay when compared with those who are not allowed to eat.
- Nutritional risk score (NRS) is validated for surgical patients and should be performed at least 10 days before surgery.
- The preoperative period should be used for conditioning regimens like prehabilitation.
- High blood glucose concentrations in patients who were normoglycemic previously are associated with increased postoperative complications.
- Excess 0.9% saline is detrimental in the perioperative period and salt and water overload of >2.5 L is associated with adverse outcome.
- Enhanced Recovery After Surgery principles are appropriate for all patients, but good results are dependent on a challenging inter-disciplinary cooperation to ensure high compliance rates.
- Inflammation is a marker for surgical complications and CRP profiling is useful.
- The effects of nutrients are dissociated from nutrition and there is a role for pharmaconutrition.
- Dysbiosis contributes to inflammation – the effects of pre-, pro- and synbiotics depends on species, strains and adjuncts.
- Postoperative fatigue inhibits voluntary exercise, immobilization induces anabolic resistance, and the lower the anabolic response to feeding, the higher the muscle loss.
- Perioperative nursing in the hospital and community after discharge is a key component for good outcome.
- A strong partnership between the surgeon and patient/family/caregivers is needed to overcome complicated postoperative courses.

concepts which are necessary for increasing implementation which include promotion of education including resources such as the ESPEN journals, ESPEN consensus papers, the life long learning (LLL) courses and live-expert courses, as well as improved communication between members of the multidisciplinary team. This may be facilitated by the creation of specialty-specific guidelines including a simplified version for community-based care as well as a patient-orientated version.

20. Conclusions

These proceedings of the ESPEN Symposium on perioperative care encompass the scientific basis of nutritional and metabolic care in the perioperative period and also suggest areas for future research and change in policy. The main take-home messages are summarized in [Table 4](#).

Funding

This symposium was supported by a grant from the European Society for Clinical Nutrition and Metabolism (ESPEN), Luxembourg.

Conflict of Interest

LG, AA, RB, KD, PLG, DHJ, SK, ZK, DCM, KER, MPS, AS and RS: None to declare.

DNL: Unrestricted grant from B. Braun and speaker's honoraria from B. Braun, Baxter Healthcare, Fresenius Kabi and Shire for unrelated work.

NEPD: Consultancy, grant and speaker's honoraria from Abbott Nutrition for unrelated work.

MH: Unrestricted grants from Abbott Nutrition and Fresenius Kabi for unrelated work.

OL: Consultancy fees from Nutricia, Pharmacosmos, Enhanced Medical Nutrition and Merck. Speaker's honoraria from Fresenius Kabi, Nutricia, Encare AB and B. Braun for unrelated work. He is also the Chairman of the ERAS Society and founder and shareholder of Encare AB.

RJES: Grant from Novartis and speaker's honoraria from Helsinn for unrelated work.

ZS: Grants from Nestle Health Science, Fresenius Kabi, Abbott Nutrition and Baxter Healthcare for unrelated work.

AW: Grants from Baxter Healthcare and Danone for unrelated work. Speaker's honoraria from Baxter, B. Braun, Ethicon, Falk Foundation and Fresenius Kabi.

Author contributions

All authors participated in the Symposium and were involved with the writing of the manuscript, critical revision and final approval.

Acknowledgements

The authors dedicate this manuscript to the memory of Mr. Ralph Stockley who sadly passed away shortly after the Symposium. Ralph was the ideal patient: stoic in adversity, yet understanding of the complexities of surgical care and always ready to provide both appreciation and constructive criticism. He will be missed greatly.

References

- [1] Weimann A, Braga M, Carli F, Higashiguchi T, Hubner M, Klek S, et al. ESPEN guideline: clinical nutrition in surgery. *Clin Nutr* 2017;36:623–50.
- [2] Chernoff R. An overview of tube feeding: from ancient times to the future. *Nutr Clin Pract* 2006;21:408–10.
- [3] Dudrick SJ. History of parenteral nutrition. *J Am Coll Nutr* 2009;28:243–51.
- [4] Harkness L. The history of enteral nutrition therapy: from raw eggs and nasal tubes to purified amino acids and early postoperative jejunal delivery. *J Am Diet Assoc* 2002;102:399–404.
- [5] Dukes C. A simple mode of feeding some patients by the nose. *Lancet* 1876;108:394–5.
- [6] Lind J. A treatise of the scurvy. In three parts. Containing an inquiry into the nature, causes, and cure, of that disease, together with a critical and chronological view of what has been published on the subject. Edinburgh: A. Kincaid & A. Donaldson; 1753.
- [7] Mozaffarian D, Rosenberg I, Uauy R. History of modern nutrition science—implications for current research, dietary guidelines, and food policy. *BMJ* 2018;361:k2392.
- [8] Studley HO. Prognosis of weight loss. *JAMA* 1936;106:458–60.
- [9] Fearon K, Strasser F, Anker SD, Bosaeus I, Bruera E, Fainsinger RL, et al. Definition and classification of cancer cachexia: an international consensus. *Lancet Oncol* 2011;12:489–95.
- [10] Morley JE, Abbatecola AM, Argiles JM, Baracos V, Bauer J, Bhasin S, et al. Sarcopenia with limited mobility: an international consensus. *J Am Med Dir Assoc* 2011;12:403–9.
- [11] Arends J, Baracos V, Bertz H, Bozzetti F, Calder PC, Deutz NEP, et al. ESPEN expert group recommendations for action against cancer-related malnutrition. *Clin Nutr* 2017;36:1187–96.
- [12] Baracos VE, Martin L, Korc M, Guttridge DC, Fearon KCH. Cancer-associated cachexia. *Nat Rev Dis Primers* 2018;4:17105.
- [13] Laird B, Fallon M. Treating cancer cachexia: an evolving landscape. *Ann Oncol* 2017;28:2055–6.
- [14] Diakos CI, Charles KA, McMillan DC, Clarke SJ. Cancer-related inflammation and treatment effectiveness. *Lancet Oncol* 2014;15:e493–503.
- [15] Cederholm T, Barazzoni R, Austin P, Ballmer P, Biolo G, Bischoff SC, et al. ESPEN guidelines on definitions and terminology of clinical nutrition. *Clin Nutr* 2017;36:49–64.
- [16] Cederholm T, Jensen GL, Correia M, Gonzalez MC, Fukushima R, Higashiguchi T, et al. GLIM criteria for the diagnosis of malnutrition – a

- consensus report from the global clinical nutrition community. *Clin Nutr* 2019;38:1–9.
- [17] Stratton RJ, Hackston A, Longmore D, Dixon R, Price S, Stroud M, et al. Malnutrition in hospital outpatients and inpatients: prevalence, concurrent validity and ease of use of the 'malnutrition universal screening tool' ('MUST') for adults. *Br J Nutr* 2004;92:799–808.
- [18] Almasaudi AS, McSorley ST, Dolan RD, Edwards CA, McMillan DC. The relation between malnutrition universal screening tool (MUST), computed tomography-derived body composition, systemic inflammation, and clinical outcomes in patients undergoing surgery for colorectal cancer. *Am J Clin Nutr* 2019;110:1327–34.
- [19] Miller J, Wells L, Nwulu U, Currow D, Johnson MJ, Skipworth RJE. Validated screening tools for the assessment of cachexia, sarcopenia, and malnutrition: a systematic review. *Am J Clin Nutr* 2018;108:1196–208.
- [20] Fowler AJ, Abbott TEF, Prowle J, Pearse RM. Age of patients undergoing surgery. *Br J Surg* 2019;106:1012–8.
- [21] Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, et al. Sarcopenia: European consensus on definition and diagnosis: report of the European Working Group on Sarcopenia in Older People. *Age Ageing* 2010;39:412–23.
- [22] Fielding RA, Vellas B, Evans WJ, Bhasin S, Morley JE, Newman AB, et al. Sarcopenia: an undiagnosed condition in older adults. Current consensus definition: prevalence, etiology, and consequences. International Working Group on Sarcopenia. *J Am Med Dir Assoc* 2011;12:249–56.
- [23] Chen LK, Liu LK, Woo J, Assantachai P, Auyeung TW, Bahyah KS, et al. Sarcopenia in Asia: consensus report of the Asian Working Group for Sarcopenia. *J Am Med Dir Assoc* 2014;15:95–101.
- [24] Dam TT, Peters KW, Fragala M, Cawthon PM, Harris TB, McLean R, et al. An evidence-based comparison of operational criteria for the presence of sarcopenia. *J Gerontol A Biol Sci Med Sci* 2014;69:584–90.
- [25] Muscaritoli M, Anker SD, Argiles J, Aversa Z, Bauer JM, Biolo G, et al. Consensus definition of sarcopenia, cachexia and pre-cachexia: joint document elaborated by special interest groups (SIG) "cachexia-anorexia in chronic wasting diseases" and "nutrition in geriatrics". *Clin Nutr* 2010;29:154–9.
- [26] Fearon K, Arends J, Baracos V. Understanding the mechanisms and treatment options in cancer cachexia. *Nat Rev Clin Oncol* 2013;10:90–9.
- [27] Ratnayake CB, Loveday BP, Shrikhande SV, Windsor JA, Pandanaboyana S. Impact of preoperative sarcopenia on postoperative outcomes following pancreatic resection: a systematic review and meta-analysis. *Pancreatology* 2018;18:996–1004.
- [28] Kamarajah SK, Bundred J, Tan BHL. Body composition assessment and sarcopenia in patients with gastric cancer: a systematic review and meta-analysis. *Gastric Cancer* 2019;22:10–22.
- [29] Matsunaga T, Miyata H, Sugimura K, Motoori M, Asukai K, Yanagimoto Y, et al. Prognostic significance of sarcopenia and systemic inflammatory response in patients with esophageal cancer. *Anticancer Res* 2019;39:449–58.
- [30] Esser H, Resch T, Pamminger M, Mutschlechner B, Troppmair J, Riedmann M, et al. Preoperative assessment of muscle mass using computerized tomography scans to predict outcomes following orthotopic liver transplantation. *Transplantation* 2019;103:2506–14.
- [31] Nakanishi R, Oki E, Sasaki S, Hirose K, Jogo T, Eda Hiro K, et al. Sarcopenia is an independent predictor of complications after colorectal cancer surgery. *Surg Today* 2018;48:151–7.
- [32] Mintziras I, Miligkos M, Wachter S, Manoharan J, Maurer E, Bartsch DK. Sarcopenia and sarcopenic obesity are significantly associated with poorer overall survival in patients with pancreatic cancer: systematic review and meta-analysis. *Int J Surg* 2018;59:19–26.
- [33] Martin L, Birdsell L, Macdonald N, Reiman T, Clandinin MT, McCargar LJ, et al. Cancer cachexia in the age of obesity: skeletal muscle depletion is a powerful prognostic factor, independent of body mass index. *J Clin Oncol* 2013;31:1539–47.
- [34] Dabiri S, Popuri K, Cespedes Feliciano EM, Caan BJ, Baracos VE, Beg MF. Muscle segmentation in axial computed tomography (CT) images at the lumbar (L3) and thoracic (T4) levels for body composition analysis. *Comput Med Imag Graph* 2019;75:47–55.
- [35] van Dijk DPJ, Bakers FCH, Sanduleanu S, Vaes RDW, Rensen SS, Dejong CHC, et al. Myosteatosis predicts survival after surgery for periampullary cancer: a novel method using MRI. *HPB (Oxford)* 2018;20:715–20.
- [36] Ataseven B, Luengo TG, du Bois A, Waltering KU, Traut A, Heitz F, et al. Skeletal muscle attenuation (sarcopenia) predicts reduced overall survival in patients with advanced epithelial ovarian cancer undergoing primary debulking surgery. *Ann Surg Oncol* 2018;25:3372–9.
- [37] Berkel AEM, Klaase JM, de Graaff F, Brusse-Keizer MGJ, Bongers BC, van Meeteren NLU. Patient's skeletal muscle radiation attenuation and sarcopenic obesity are associated with postoperative morbidity after neoadjuvant chemoradiation and resection for rectal cancer. *Dig Surg* 2019;36:376–83.
- [38] Xiao J, Caan BJ, Weltzien E, Cespedes Feliciano EM, Kroenke CH, Meyerhardt JA, et al. Associations of pre-existing co-morbidities with skeletal muscle mass and radiodensity in patients with non-metastatic colorectal cancer. *J Cachexia Sarcopenia Muscle* 2018;9:654–63.
- [39] Skipworth RJ, Stewart GD, Dejong CH, Preston T, Fearon KC. Pathophysiology of cancer cachexia: much more than host-tumour interaction? *Clin Nutr* 2007;26:667–76.
- [40] Johns N, Stretch C, Tan BH, Solheim TS, Sorhaug S, Stephens NA, et al. New genetic signatures associated with cancer cachexia as defined by low skeletal muscle index and weight loss. *J Cachexia Sarcopenia Muscle* 2017;8:122–30.
- [41] Bozzetti F, Mariani L, Lo Vullo S, Amerio ML, Biffi R, Caccialanza G, et al. The nutritional risk in oncology: a study of 1,453 cancer outpatients. *Support Care Cancer* 2012;20:1919–28.
- [42] Evans WJ, Morley JE, Argiles J, Bales C, Baracos V, Guttridge D, et al. Cachexia: a new definition. *Clin Nutr* 2008;27:793–9.
- [43] Fearon KC, Voss AC, Hustead DS, Cancer Cachexia Study Group. Definition of cancer cachexia: effect of weight loss, reduced food intake, and systemic inflammation on functional status and prognosis. *Am J Clin Nutr* 2006;83:1345–50.
- [44] Crossland H, Skirrow S, Puthuchery ZA, Constantin-Teodosiu D, Greenhaff PL. The impact of immobilisation and inflammation on the regulation of muscle mass and insulin resistance: different routes to similar endpoints. *J Physiol* 2019;597:1259–70.
- [45] Galvan E, Arentson-Lantz E, Lamon S, Paddon-Jones D. Protecting skeletal muscle with protein and amino acid during periods of disuse. *Nutrients* 2016;8.
- [46] Deutz NE, Pereira SL, Hays NP, Oliver JS, Edens NK, Evans CM, et al. Effect of beta-hydroxy-beta-methylbutyrate (HMB) on lean body mass during 10 days of bed rest in older adults. *Clin Nutr* 2013;32:704–12.
- [47] Drummond MJ, Dickinson JM, Fry CS, Walker DK, Gundermann DM, Reidy PT, et al. Bed rest impairs skeletal muscle amino acid transporter expression, mtorc1 signaling, and protein synthesis in response to essential amino acids in older adults. *Am J Physiol Endocrinol Metab* 2012;302:E1113–22.
- [48] Hvid LG, Suetta C, Nielsen JH, Jensen MM, Frandsen U, Ortenblad N, et al. Aging impairs the recovery in mechanical muscle function following 4 days of disuse. *Exp Gerontol* 2014;52:1–8.
- [49] Ferrando AA, Tipton KD, Bamman MM, Wolfe RR. Resistance exercise maintains skeletal muscle protein synthesis during bed rest. *J Appl Physiol* (1985) 1997;82:807–10.
- [50] Bos C, Benamouzig R, Bruhat A, Roux C, Valensi P, Ferriere F, et al. Nutritional status after short-term dietary supplementation in hospitalized malnourished geriatric patients. *Clin Nutr* 2001;20:225–33.
- [51] Ferrando AA, Paddon-Jones D, Hays NP, Kortebein P, Ronsen O, Williams RH, et al. EAA supplementation to increase nitrogen intake improves muscle function during bed rest in the elderly. *Clin Nutr* 2010;29:18–23.
- [52] Paddon-Jones D, Sheffield-Moore M, Urban RJ, Sanford AP, Aarsland A, Wolfe RR, et al. Essential amino acid and carbohydrate supplementation ameliorates muscle protein loss in humans during 28 days bedrest. *J Clin Endocrinol Metab* 2004;89:4351–8.
- [53] Puthuchery ZA, Rawal J, McPhail M, Connolly B, Ratnayake G, Chan P, et al. Acute skeletal muscle wasting in critical illness. *JAMA* 2013;310:1591–600.
- [54] Williams JP, Phillips BE, Smith K, Atherton PJ, Rankin D, Selby AL, et al. Effect of tumor burden and subsequent surgical resection on skeletal muscle mass and protein turnover in colorectal cancer patients. *Am J Clin Nutr* 2012;96:1064–70.
- [55] Schindler K, Pichard C, Sulz I, Volkert D, Streicher M, Singer P, et al. nutritionDay: 10 years of growth. *Clin Nutr* 2017;36:1207–14.
- [56] Frantal S, Pernicka E, Hiesmayr M, Schindler K, Bauer P. Length bias correction in one-day cross-sectional assessments - the nutritionday study. *Clin Nutr* 2016;35:522–7.
- [57] Pirlich M, Schutz T, Norman K, Gastell S, Lubke HJ, Bischoff SC, et al. The German hospital malnutrition study. *Clin Nutr* 2006;25:563–72.
- [58] Pikul J, Sharpe MD, Lowndes R, Ghent CN. Degree of preoperative malnutrition is predictive of postoperative morbidity and mortality in liver transplant recipients. *Transplantation* 1994;57:469–72.
- [59] Meguid MM, Debonis D, Meguid V, Hill LR, Terz JJ. Complications of abdominal operations for malignant disease. *Am J Surg* 1988;156:341–5.
- [60] Kondrup J, Allison SP, Elia M, Vellas B, Plauth M, Educational, et al. ESPEN guidelines for nutrition screening 2002. *Clin Nutr* 2003;22:415–21.
- [61] Veterans Affairs Total Parenteral Nutrition Cooperative Study Group. Perioperative total parenteral nutrition in surgical patients. *N Engl J Med* 1991;325:525–32.
- [62] Detsky AS, McLaughlin JR, Baker JP, Johnston N, Whittaker S, Mendelson RA, et al. What is subjective global assessment of nutritional status? *J Parenter Enter Nutr* 1987;11:8–13.
- [63] Chohendy R, Rubenstein LZ, Eledjam JJ. The mini nutritional assessment-short form for preoperative nutritional evaluation of elderly patients. *Aging (Milano)* 2001;13:293–7.
- [64] Kondrup J, Rasmussen HH, Hamberg O, Stanga Z, Ad Hoc EWG. Nutritional risk screening (NRS 2002): a new method based on an analysis of controlled clinical trials. *Clin Nutr* 2003;22:321–36.
- [65] Guo W, Ou G, Li X, Huang J, Liu J, Wei H. Screening of the nutritional risk of patients with gastric carcinoma before operation by NRS 2002 and its relationship with postoperative results. *J Gastroenterol Hepatol* 2010;25:800–3.
- [66] Schwegler I, von Holzen A, Gutzwiller JP, Schlumpf R, Muhlebach S, Stanga Z. Nutritional risk is a clinical predictor of postoperative mortality and morbidity in surgery for colorectal cancer. *Br J Surg* 2010;97:92–7.
- [67] Schiesser M, Muller S, Kirchhoff P, Breitenstein S, Schafer M, Clavien PA. Assessment of a novel screening score for nutritional risk in predicting complications in gastro-intestinal surgery. *Clin Nutr* 2008;27:565–70.
- [68] Schiesser M, Kirchhoff P, Muller MK, Schafer M, Clavien PA. The correlation of nutrition risk index, nutrition risk score, and bioimpedance analysis with

- postoperative complications in patients undergoing gastrointestinal surgery. *Surgery* 2009;145:519–26.
- [69] Sun Z, Kong XJ, Jing X, Deng RJ, Tian ZB. Nutritional risk screening 2002 as a predictor of postoperative outcomes in patients undergoing abdominal surgery: a systematic review and meta-analysis of prospective cohort studies. *PLoS One* 2015;10:e0132857.
- [70] Reber E, Gomes F, Vasiloglou MF, Schuetz P, Stanga Z. Nutritional risk screening and assessment. *J Clin Med* 2019;8.
- [71] Fearon KC, Ljungqvist O, Von Meyenfeldt M, Revhaug A, Dejong CH, Lassen K, et al. Enhanced Recovery after Surgery: a consensus review of clinical care for patients undergoing colonic resection. *Clin Nutr* 2005;24:466–77.
- [72] Horowitz M, Neeman E, Sharon E, Ben-Eliyahu S. Exploiting the critical perioperative period to improve long-term cancer outcomes. *Nat Rev Clin Oncol* 2015;12:213–26.
- [73] Allison SP. The uses and limitations of nutritional support the arvid wretling lecture given at the 14th ESPEN Congress in Vienna, 1992. *Clin Nutr* 1992;11:319–30.
- [74] Arends J, Bachmann P, Baracos V, Barthelemy N, Bertz H, Bozzetti F, et al. ESPEN guidelines on nutrition in cancer patients. *Clin Nutr* 2017;36:11–48.
- [75] Gould DW, Lahart I, Carmichael AR, Koutedakis Y, Metsios GS. Cancer cachexia prevention via physical exercise: molecular mechanisms. *J Cachexia Sarcopenia Muscle* 2013;4:111–24.
- [76] Bauer J, Biolo G, Cederholm T, Cesari M, Cruz-Jentoft AJ, Morley JE, et al. Evidence-based recommendations for optimal dietary protein intake in older people: a position paper from the PROT-AGE study group. *J Am Med Dir Assoc* 2013;14:542–59.
- [77] Elia M, Normand C, Norman K, Laviano A. A systematic review of the cost and cost effectiveness of using standard oral nutritional supplements in the hospital setting. *Clin Nutr* 2016;35:370–80.
- [78] Fukuda Y, Yamamoto K, Hirao M, Nishikawa K, Maeda S, Haraguchi N, et al. Prevalence of malnutrition among gastric cancer patients undergoing gastrectomy and optimal preoperative nutritional support for preventing surgical site infections. *Ann Surg Oncol* 2015;22(Suppl 3):S778–85.
- [79] Awad S, Varadhan KK, Ljungqvist O, Lobo DN. A meta-analysis of randomised controlled trials on preoperative oral carbohydrate treatment in elective surgery. *Clin Nutr* 2013;32:34–44.
- [80] Amer MA, Smith MD, Herbison GP, Plank LD, McCall JL. Network meta-analysis of the effect of preoperative carbohydrate loading on recovery after elective surgery. *Br J Surg* 2017;104:187–97.
- [81] Gianotti L, Biffi R, Sandini M, Marrelli D, Vignali A, Caccialanza R, et al. Preoperative oral carbohydrate load versus placebo in major elective abdominal surgery (PROCY): a randomized, placebo-controlled, multicenter, phase III trial. *Ann Surg* 2018;267:623–30.
- [82] Minnella EM, Bousquet-Dion G, Awasthi R, Scheede-Bergdahl C, Carli F. Multimodal prehabilitation improves functional capacity before and after colorectal surgery for cancer: a five-year research experience. *Acta Oncol* 2017;56:295–300.
- [83] Gillis C, Buhler K, Bresee L, Carli F, Gramlich L, Culos-Reed N, et al. Effects of nutritional prehabilitation, with and without exercise, on outcomes of patients who undergo colorectal surgery: a systematic review and meta-analysis. *Gastroenterology* 2018;155:391–410 e4.
- [84] Gillis C, Fenton TR, Sajobi TT, Minnella EM, Awasthi R, Loiselle SE, et al. Trimodal prehabilitation for colorectal surgery attenuates post-surgical losses in lean body mass: a pooled analysis of randomized controlled trials. *Clin Nutr* 2019;38:1053–60.
- [85] Dunne DF, Jack S, Jones RP, Jones L, Lythgoe DT, Malik HZ, et al. Randomized clinical trial of prehabilitation before planned liver resection. *Br J Surg* 2016;103:504–12.
- [86] Yamamoto K, Nagatsuma Y, Fukuda Y, Hirao M, Nishikawa K, Miyamoto A, et al. Effectiveness of a preoperative exercise and nutritional support program for elderly sarcopenic patients with gastric cancer. *Gastric Cancer* 2017;20:913–8.
- [87] Barberan-Garcia A, Ubre M, Roca J, Lacy AM, Burgos F, Risco R, et al. Personalised prehabilitation in high-risk patients undergoing elective major abdominal surgery: a randomized blinded controlled trial. *Ann Surg* 2018;267:50–6.
- [88] Dhatriya K, Flanagan D, Hilton L, Kilvert A, Levy N, Rayman G, et al. Management of adults with diabetes undergoing surgery and elective procedures: improving standards. London: Diabetes UK; 2011. Available at: www.diabetologists-abcd.org.uk/JBDS/JBDS_IP_Surgery_Adults_Full.pdf. [Accessed 10 March 2020].
- [89] Pournaras DJ, Photi ES, Barnett N, Challand CP, Chatzizacharias NA, Dlamini NP, et al. Assessing the quality of primary care referrals to surgery of patients with diabetes in the east of England: a multi-centre cross-sectional cohort study. *Int J Clin Pract* 2017;71:e12971.
- [90] Frisch A, Chandra P, Smiley D, Peng L, Rizzo M, Gatcliffe C, et al. Prevalence and clinical outcome of hyperglycemia in the perioperative period in noncardiac surgery. *Diabetes Care* 2010;33:1783–8.
- [91] Kotagal M, Symons RC, Hirsch IB, Umpierrez GE, Dellinger EP, Farrokhi ET, et al. Perioperative hyperglycemia and risk of adverse events among patients with and without diabetes. *Ann Surg* 2015;261:97–103.
- [92] Kwon S, Thompson R, Dellinger P, Yanez D, Farrokhi E, Flum D. Importance of perioperative glycemic control in general surgery: a report from the surgical care and outcomes assessment program. *Ann Surg* 2013;257:8–14.
- [93] Critchley JA, Carey IM, Harris T, DeWilde S, Hosking FJ, Cook DG. Glycemic control and risk of infections among people with type 1 or type 2 diabetes in a large primary care cohort study. *Diabetes Care* 2018;41:2127–35.
- [94] Jones CE, Graham LA, Morris MS, Richman JS, Hollis RH, Wahl TS, et al. Association between preoperative hemoglobin A1c levels, postoperative hyperglycemia, and readmissions following gastrointestinal surgery. *JAMA Surg* 2017;152:1031–8.
- [95] National Collaborating Centre for Acute Care (UK). Preoperative tests. The use of routine preoperative tests for elective surgery. London: National Collaborating Centre for Acute Care (UK); 2003. Available at: https://www.ncbi.nlm.nih.gov/books/NBK48489/pdf/Bookshelf_NBK48489.pdf. [Accessed 10 March 2020].
- [96] Dhatriya KK, Wiles MD. Pre-operative testing guidelines: a NICE try but not enough. *Anaesthesia* 2016;71:1403–7.
- [97] Lobo DN. Fluid, electrolytes and nutrition: physiological and clinical aspects. *Proc Nutr Soc* 2004;63:453–66.
- [98] Keys A, Brozek J, Henschel A, Mickelsen O, Taylor HF. The biology of human starvation. Minneapolis: University of Minnesota Press; 1950.
- [99] Moore FD. Metabolic care of the surgical patient. Philadelphia: W. B. Saunders; 1959.
- [100] Shizgal HM. The effect of malnutrition on body composition. *Surg Gynecol Obstet* 1981;152:22–6.
- [101] Winick M. Hunger disease: studies by the Jewish physicians in the Warsaw ghetto. New York: Wiley; 1979.
- [102] Wilkinson AW, Nagy G, Billing BH, Stewart CP. Excretion of chloride and sodium after surgical operations. *Lancet* 1949;1:640–4.
- [103] Wilkinson AW, Billing BH, Nagy G, Stewart CP. Excretion of potassium after partial gastrectomy. *Lancet* 1950;2:135–7.
- [104] Thiele RH, Raghunathan K, Brudney CS, Lobo DN, Martin D, Senagore A, et al. American Society for Enhanced Recovery (ASER) and Perioperative Quality Initiative (POQI) joint consensus statement on perioperative fluid management within an enhanced recovery pathway for colorectal surgery. *Perioper Med (Lond)* 2016;5:24.
- [105] Gan TJ, Soppitt A, Maroof M, el-Moalem H, Robertson KM, Moretti E, et al. Goal-directed intraoperative fluid administration reduces length of hospital stay after major surgery. *Anesthesiology* 2002;97:820–6.
- [106] Noblett SE, Snowden CP, Shenton BK, Horgan AF. Randomized clinical trial assessing the effect of Doppler-optimized fluid management on outcome after elective colorectal resection. *Br J Surg* 2006;93:1069–76.
- [107] Feldheiser A, Aziz O, Baldini G, Cox BP, Fearon KC, Feldman LS, et al. Enhanced Recovery after Surgery (ERAS) for gastrointestinal surgery, part 2: consensus statement for anaesthesia practice. *Acta Anaesthesiol Scand* 2016;60:289–334.
- [108] Rollins KE, Lobo DN. Intraoperative goal-directed fluid therapy in elective major abdominal surgery: a meta-analysis of randomized controlled trials. *Ann Surg* 2016;263:465–76.
- [109] Rollins KE, Mathias NC, Lobo DN. Meta-analysis of goal-directed fluid therapy using transoesophageal Doppler monitoring in patients undergoing elective colorectal surgery. *BJS Open* 2019;3:606–16.
- [110] Lassen K, Kjaeve J, Fetveit T, Trano G, Sigurdsson HK, Horn A, et al. Allowing normal food at will after major upper gastrointestinal surgery does not increase morbidity: a randomized multicenter trial. *Ann Surg* 2008;247:721–9.
- [111] Zhuang CL, Ye XZ, Zhang CJ, Dong QT, Chen BC, Yu Z. Early versus traditional postoperative oral feeding in patients undergoing elective colorectal surgery: a meta-analysis of randomized clinical trials. *Dig Surg* 2013;30:225–32.
- [112] NICE Guideline [CG174]. Intravenous fluid therapy for adults in hospital. London: National Institute for Health and Care Excellence; 2013. Available at: <https://www.nice.org.uk/guidance/cg174>. [Accessed 10 March 2020].
- [113] Lobo DN, Bostock KA, Neal KR, Perkins AC, Rowlands BJ, Allison SP. Effect of salt and water balance on recovery of gastrointestinal function after elective colonic resection: a randomised controlled trial. *Lancet* 2002;359:1812–8.
- [114] Van Regenmortel N, De Weerd T, Van Craenenbroeck AH, Roelant E, Verbrugge W, Dams K, et al. Effect of isotonic versus hypotonic maintenance fluid therapy on urine output, fluid balance, and electrolyte homeostasis: a crossover study in fasting adult volunteers. *Br J Anaesth* 2017;118:892–900.
- [115] Gould TH, Grace K, Thorne G, Thomas M. Effect of thoracic epidural anaesthesia on colonic blood flow. *Br J Anaesth* 2002;89:446–51.
- [116] Holte K, Foss NB, Svensen C, Lund C, Madsen JL, Kehlet H. Epidural anesthesia, hypotension, and changes in intravascular volume. *Anesthesiology* 2004;100:281–6.
- [117] Lobo DN. Fluid overload and surgical outcome: another piece in the jigsaw. *Ann Surg* 2009;249:186–8.
- [118] Shin CH, Long DR, McLean D, Grabitz SD, Ladha K, Timm FP, et al. Effects of intraoperative fluid management on postoperative outcomes: a hospital registry study. *Ann Surg* 2018;267:1084–92.
- [119] Thacker JK, Mountford WK, Ernst FR, Krukus MR, Mythen MM. Perioperative fluid utilization variability and association with outcomes: considerations for enhanced recovery efforts in sample US surgical populations. *Ann Surg* 2016;263:502–10.
- [120] Varadhan KK, Lobo DN. A meta-analysis of randomised controlled trials of intravenous fluid therapy in major elective open abdominal surgery: getting the balance right. *Proc Nutr Soc* 2010;69:488–98.

- [121] Lobo DN, Stanga Z, Aloysius MM, Wicks C, Nunes QM, Ingram KL, et al. Effect of volume loading with 1 liter intravenous infusions of 0.9% saline, 4% succinylated gelatine (gelofusine) and 6% hydroxyethyl starch (voluven) on blood volume and endocrine responses: a randomized, three-way crossover study in healthy volunteers. *Crit Care Med* 2010;38:464–70.
- [122] Drummer C, Gerzer R, Heer M, Molz B, Bie P, Schlossberger M, et al. Effects of an acute saline infusion on fluid and electrolyte metabolism in humans. *Am J Physiol* 1992;262:F744–54.
- [123] Lobo DN, Stanga Z, Simpson JA, Anderson JA, Rowlands BJ, Allison SP. Dilution and redistribution effects of rapid 2-litre infusions of 0.9% (w/v) saline and 5% (w/v) dextrose on haematological parameters and serum biochemistry in normal subjects: a double-blind crossover study. *Clin Sci (Lond)* 2001;101:173–9.
- [124] Reid F, Lobo DN, Williams RN, Rowlands BJ, Allison SP. (ab)normal saline and physiological Hartmann's solution: a randomized double-blind crossover study. *Clin Sci (Lond)* 2003;104:17–24.
- [125] Wilkes NJ, Woolf R, Mutch M, Mallett SV, Peachey T, Stephens R, et al. The effects of balanced versus saline-based hetastarch and crystalloid solutions on acid-base and electrolyte status and gastric mucosal perfusion in elderly surgical patients. *Anesth Analg* 2001;93:811–6.
- [126] Williams EL, Hildebrand KL, McCormick SA, Bedel MJ. The effect of intravenous lactated ringer's solution versus 0.9% sodium chloride solution on serum osmolality in human volunteers [see comments]. *Anesth Analg* 1999;88:999–1003.
- [127] Chowdhury AH, Lobo DN. Fluids and gastrointestinal function. *Curr Opin Clin Nutr Metab Care* 2011;14:469–76.
- [128] Lobo DN, Allison SP. Fluid, electrolyte and nutrient replacement. In: Burnand KG, Young AE, Lucas J, Rowlands BJ, Scholefield J, editors. *The new Aird's companion in surgical studies*. 3rd ed. London: Churchill Livingstone; 2005. p. 20–41.
- [129] Friedman J, Lussiez A, Sullivan J, Wang S, Englesbe M. Implications of sarcopenia in major surgery. *Nutr Clin Pract* 2015;30:175–9.
- [130] Muckart DJ, Bhagwanjee S. American College of Chest Physicians/Society of Critical Care Medicine Consensus Conference definitions of the systemic inflammatory response syndrome and allied disorders in relation to critically injured patients. *Crit Care Med* 1997;25:1789–95.
- [131] Gabay C, Kushner I. Acute-phase proteins and other systemic responses to inflammation. *N Engl J Med* 1999;340:448–54.
- [132] Watt DG, Horgan PG, McMillan DC. Routine clinical markers of the magnitude of the systemic inflammatory response after elective operation: a systematic review. *Surgery* 2015;157:362–80.
- [133] Platt JJ, Ramanathan ML, Crosbie RA, Anderson JH, McKee RF, Horgan PG, et al. C-reactive protein as a predictor of postoperative infective complications after curative resection in patients with colorectal cancer. *Ann Surg Oncol* 2012;19:4168–77.
- [134] Gans SL, Atema JJ, van Dieren S, Groot Koerkamp B, Boermeester MA. Diagnostic value of C-reactive protein to rule out infectious complications after major abdominal surgery: a systematic review and meta-analysis. *Int J Colorectal Dis* 2015;30:861–73.
- [135] Straatman J, Harmsen AM, Cuesta MA, Berkhof J, Jansma EP, van der Peet DL. Predictive value of C-reactive protein for major complications after major abdominal surgery: a systematic review and pooled-analysis. *PLoS One* 2015;10:e0132995.
- [136] Watt DG, McSorley ST, Horgan PG, McMillan DC. Enhanced recovery after surgery: which components, if any, impact on the systemic inflammatory response following colorectal surgery?: a systematic review. *Medicine (Baltim)* 2015;94:e1286.
- [137] McSorley ST, Horgan PG, McMillan DC. The impact of preoperative corticosteroids on the systemic inflammatory response and postoperative complications following surgery for gastrointestinal cancer: a systematic review and meta-analysis. *Crit Rev Oncol Hematol* 2016;101:139–50.
- [138] Golder AM, Steele CW, Conn D, MacKay GJ, McMillan DC, Horgan PG, et al. Effect of preoperative oral antibiotics in combination with mechanical bowel preparation on inflammatory response and short-term outcomes following left-sided colonic and rectal resections. *BJS Open* 2019;3:830–9.
- [139] Rollins KE, Javanmard-Emamghissi H, Acheson AG, Lobo DN. The role of oral antibiotic preparation in elective colorectal surgery: a meta-analysis. *Ann Surg* 2019;270:43–58.
- [140] Ljungqvist O, Scott M, Fearon KC. Enhanced recovery after surgery: a review. *JAMA Surg* 2017;152:292–8.
- [141] Rollins KE, Javanmard-Emamghissi H, Lobo DN. Impact of mechanical bowel preparation in elective colorectal surgery: a meta-analysis. *World J Gastroenterol* 2018;24:519–36.
- [142] Maessen J, Dejong CH, Hausel J, Nygren J, Lassen K, Andersen J, et al. A protocol is not enough to implement an enhanced recovery programme for colorectal resection. *Br J Surg* 2007;94:224–31.
- [143] Varadhan KK, Neal KR, Dejong CH, Fearon KC, Ljungqvist O, Lobo DN. The enhanced recovery after surgery (ERAS) pathway for patients undergoing major elective open colorectal surgery: a meta-analysis of randomized controlled trials. *Clin Nutr* 2010;29:434–40.
- [144] Visioni A, Shah R, Gabriel E, Attwood K, Kukar M, Nurkin S. Enhanced recovery after surgery for noncolorectal surgery?: a systematic review and meta-analysis of major abdominal surgery. *Ann Surg* 2018;267:57–65.
- [145] Jia Y, Jin G, Guo S, Gu B, Jin Z, Gao X, et al. Fast-track surgery decreases the incidence of postoperative delirium and other complications in elderly patients with colorectal carcinoma. *Langenbeck's Arch Surg* 2014;399:77–84.
- [146] Jorgensen CC, Madsbad S, Kehlet H, Lundbeck Foundation Centre for Fast-track H, Knee Replacement Collaborative G. Postoperative morbidity and mortality in type-2 diabetics after fast-track primary total hip and knee arthroplasty. *Anesth Analg* 2015;120:230–8.
- [147] Hendry PO, Hausel J, Nygren J, Lassen K, Dejong CH, Ljungqvist O, et al. Determinants of outcome after colorectal resection within an enhanced recovery programme. *Br J Surg* 2009;96:197–205.
- [148] Gianotti L, Sandini M, Romagnoli S, Carli F, Ljungqvist O. Enhanced recovery programs in gastrointestinal surgery: actions to promote optimal perioperative nutritional and metabolic care. *Clin Nutr* 2019. <https://doi.org/10.1016/j.clnu.2019.10.023> [Epub ahead of print].
- [149] Pedziwiatr M, Pisarska M, Kisielewski M, Matlok M, Major P, Wierdak M, et al. Is ERAS in laparoscopic surgery for colorectal cancer changing risk factors for delayed recovery? *Med Oncol* 2016;33:25.
- [150] Ljungqvist O, Jonathan E. Rhoads lecture 2011: insulin resistance and enhanced recovery after surgery. *J Parenter Enter Nutr* 2012;36:389–98.
- [151] Gustafsson UO, Hausel J, Thorell A, Ljungqvist O, Soop M, Nygren J, et al. Adherence to the enhanced recovery after surgery protocol and outcomes after colorectal cancer surgery. *Arch Surg* 2011;146:571–7.
- [152] ERAS Compliance Group. The impact of enhanced recovery protocol compliance on elective colorectal cancer resection: results from an international registry. *Ann Surg* 2015;261:1153–9.
- [153] Gustafsson UO, Opielstrup H, Thorell A, Nygren J, Ljungqvist O. Adherence to the ERAS protocol is associated with 5-year survival after colorectal cancer surgery: a retrospective cohort study. *World J Surg* 2016;40:1741–7.
- [154] The Lancet Commission on Global Surgery. <https://www.lancetglobalsurgery.org/> [Accessed 10 March 2020].
- [155] Nationella Kvalitetsregister. <http://www.kvalitetsregister.se/hittaregister/registerarkiv/tjockochandtarmscancer.2315.html>.
- [156] Healy MA, Regenbogen SE, Kanters AE, Suwanabol PA, Varban OA, Campbell Jr DA, et al. Surgeon variation in complications with minimally invasive and open colectomy: results from the Michigan surgical quality collaborative. *JAMA Surg* 2017;152:860–7.
- [157] Lilot M, Ehrenfeld JM, Lee C, Harrington B, Cannesson M, Rinehart J. Variability in practice and factors predictive of total crystalloid administration during abdominal surgery: retrospective two-centre analysis. *Br J Anaesth* 2015;114:767–76.
- [158] Ljungqvist O, Thanh NX, Nelson G. ERAS-value based surgery. *J Surg Oncol* 2017;116:608–12.
- [159] Burch J, Taylor C. Patients' need for nursing telephone follow-up after enhanced recovery. *Gastrointest Nurs* 2012;10:51–8.
- [160] McShane C, Honeysett A. What is the preadmission role in preparation of patients for fast track surgery? *Aust Nurs J* 2013;20:37–9.
- [161] Gerritsen A, Besselink MG, Gouma DJ, Steenhagen E, Borel Rinkes IH, Molenaar IQ. Systematic review of five feeding routes after pancreaticoduodenectomy. *Br J Surg* 2013;100:589–98. discussion 99.
- [162] Adiamah A, Ranat R, Gomez D. Enteral versus parenteral nutrition following pancreaticoduodenectomy: a systematic review and meta-analysis. *HPB (Oxford)* 2019;21:793–801.
- [163] Lee SH, Jang JY, Kim HW, Jung MJ, Lee JG. Effects of early enteral nutrition on patients after emergency gastrointestinal surgery: a propensity score matching analysis. *Medicine (Baltim)* 2014;93:e323.
- [164] Andersen HK, Lewis SJ, Thomas S. Early enteral nutrition within 24h of colorectal surgery versus later commencement of feeding for postoperative complications. *Cochrane Database Syst Rev* 2006;4:CD004080.
- [165] Herbert G, Perry R, Andersen HK, Atkinson C, Penfold C, Lewis SJ, et al. Early enteral nutrition within 24 hours of lower gastrointestinal surgery versus later commencement for length of hospital stay and postoperative complications. *Cochrane Database Syst Rev* 2018;10:CD004080.
- [166] Willcutts KF, Chung MC, Erenberg CL, Finn KL, Schirmer BD, Byham-Gray LD. Early oral feeding as compared with traditional timing of oral feeding after upper gastrointestinal surgery: a systematic review and meta-analysis. *Ann Surg* 2016;264:54–63.
- [167] Christensen T, Kehlet H. Postoperative fatigue and changes in nutritional status. *Br J Surg* 1984;71:473–6.
- [168] Latimer LE, Constantin D, Greening NJ, Calvert L, Menon MK, Steiner MC, et al. Impact of transcutaneous neuromuscular electrical stimulation or resistance exercise on skeletal muscle mRNA expression in COPD. *Int J Chron Obstruct Pulmon Dis* 2019;14:1355–64.
- [169] Suetta C, Hvid LG, Justesen L, Christensen U, Neergaard K, Simonsen L, et al. Effects of aging on human skeletal muscle after immobilization and retraining. *J Appl Physiol* (1985) 2009;107:1172–80.
- [170] Constantin D, Menon MK, Houchen-Wolloff L, Morgan MD, Singh SJ, Greenhaff P, et al. Skeletal muscle molecular responses to resistance training and dietary supplementation in COPD. *Thorax* 2013;68:625–33.
- [171] Soeters PB, Grecu I. Have we enough glutamine and how does it work? A clinician's view. *Ann Nutr Metab* 2012;60:17–26.
- [172] Petersson B, Vinnars E, Waller SO, Wernerman J. Long-term changes in muscle free amino acid levels after elective abdominal surgery. *Br J Surg* 1992;79:212–6.

- [173] Souba WW, Klimberg VS, Plumley DA, Salloum RM, Flynn TC, Bland KI, et al. The role of glutamine in maintaining a healthy gut and supporting the metabolic response to injury and infection. *J Surg Res* 1990;48:383–91.
- [174] Zheng Y, Li F, Qi B, Luo B, Sun H, Liu S, et al. Application of perioperative immunonutrition for gastrointestinal surgery: a meta-analysis of randomized controlled trials. *Asia Pac J Clin Nutr* 2007;16(Suppl 1):253–7.
- [175] Gianotti L, Braga M, Biffi R, Bozzetti F, Mariani L. Glutamaltaly Research Group of the Italian Society of Parenteral, and Enteral Nutrition. Perioperative intravenous glutamine supplementation in major abdominal surgery for cancer: a randomized multicenter trial. *Ann Surg* 2009;250:684–90.
- [176] Ziegler TR, May AK, Hebbard G, Easley KA, Griffith DP, Dave N, et al. Efficacy and safety of glutamine-supplemented parenteral nutrition in surgical ICU patients: an American multicenter randomized controlled trial. *Ann Surg* 2016;263:646–55.
- [177] Sandini M, Nespoli L, Oldani M, Bernasconi DP, Gianotti L. Effect of glutamine dipeptide supplementation on primary outcomes for elective major surgery: systematic review and meta-analysis. *Nutrients* 2015;7:481–99.
- [178] Mayer K, Seeger W, Grimminger F. Clinical use of lipids to control inflammatory disease. *Curr Opin Clin Nutr Metab Care* 1998;1:179–84.
- [179] Wichmann MW, Thul P, Czarnetzki HD, Morlion BJ, Kemen M, Jauch KW. Evaluation of clinical safety and beneficial effects of a fish oil containing lipid emulsion (Lipoplus, MLF541): data from a prospective, randomized, multicenter trial. *Crit Care Med* 2007;35:700–6.
- [180] Pradelli L, Mayer K, Klek S, Omar Alsaleh AJ, Clark RAC, Rosenthal MD, et al. Omega-3 fatty-acid enriched parenteral nutrition in hospitalized patients: systematic review with meta-analysis and trial sequential analysis. *J Parenter Enter Nutr* 2020;44:44–57.
- [181] Li NN, Zhou Y, Qin XP, Chen Y, He D, Feng JY, et al. Does intravenous fish oil benefit patients post-surgery? A meta-analysis of randomised controlled trials. *Clin Nutr* 2014;33:226–39.
- [182] Adiamah A, Skorepa P, Weimann A, Lobo DN. The impact of preoperative immune modulating nutrition on outcomes in patients undergoing surgery for gastrointestinal cancer: a systematic review and meta-analysis. *Ann Surg* 2019;270:247–56.
- [183] Marimuthu K, Varadhan KK, Ljungqvist O, Lobo DN. A meta-analysis of the effect of combinations of immune modulating nutrients on outcome in patients undergoing major open gastrointestinal surgery. *Ann Surg* 2012;255:1060–8.
- [184] Moya P, Miranda E, Soriano-Irigaray L, Arroyo A, Aguilar MD, Bellon M, et al. Perioperative immunonutrition in normo-nourished patients undergoing laparoscopic colorectal resection. *Surg Endosc* 2016;30:4946–53.
- [185] WHO/FAO. Health and nutritional properties of probiotics in food including powder milk with live lactic acid bacteria; report of a joint FAO/WHO expert consultation on evaluation of health and nutritional properties of probiotics in food including powder milk with live lactic acid bacteria. Rome: Food and Agriculture Organization of the United Nations, World Health Organization; 2001.
- [186] Slavin J. Fiber and prebiotics: mechanisms and health benefits. *Nutrients* 2013;5:1417–35.
- [187] Allen SJ, Martinez EG, Gregorio GV, Dans LF. Probiotics for treating acute infectious diarrhoea. *Cochrane Database Syst Rev* 2010;11:CD003048.
- [188] Huang JS, Bousvaros A, Lee JW, Diaz A, Davidson EJ. Efficacy of probiotic use in acute diarrhea in children: a meta-analysis. *Dig Dis Sci* 2002;47:2625–34.
- [189] Szajewska H, Mrukowicz JZ. Probiotics in the treatment and prevention of acute infectious diarrhea in infants and children: a systematic review of published randomized, double-blind, placebo-controlled trials. *J Pediatr Gastroenterol Nutr* 2001;33(Suppl 2):S17–25.
- [190] Van Niel CW, Feudtner C, Garrison MM, Christakis DA. Lactobacillus therapy for acute infectious diarrhea in children: a meta-analysis. *Pediatrics* 2002;109:678–84.
- [191] McFarland LV. Meta-analysis of probiotics for the prevention of traveler's diarrhea. *Trav Med Infect Dis* 2007;5:97–105.
- [192] Johnston BC, Supina AL, Ospina M, Vohra S. Probiotics for the prevention of pediatric antibiotic-associated diarrhea. *Cochrane Database Syst Rev* 2007;2:CD004827.
- [193] Johnston BC, Supina AL, Vohra S. Probiotics for pediatric antibiotic-associated diarrhea: a meta-analysis of randomized placebo-controlled trials. *CMAJ* 2006;175:377–83.
- [194] Szajewska H, Rusczyński M, Radzikowski A. Probiotics in the prevention of antibiotic-associated diarrhea in children: a meta-analysis of randomized controlled trials. *J Pediatr* 2006;149:367–72.
- [195] Cremonini F, Di Caro S, Nista EC, Bartolozzi F, Capelli G, Gasbarrini G, et al. Meta-analysis: the effect of probiotic administration on antibiotic-associated diarrhoea. *Aliment Pharmacol Ther* 2002;16:1461–7.
- [196] D'Souza AL, Rajkumar C, Cooke J, Bulpitt CJ. Probiotics in prevention of antibiotic associated diarrhoea: meta-analysis. *BMJ* 2002;324:1361.
- [197] Hawrelak JA, Whitten DL, Myers SP. Is Lactobacillus rhamnosus GG effective in preventing the onset of antibiotic-associated diarrhoea: a systematic review. *Digestion* 2005;72:51–6.
- [198] Szajewska H, Mrukowicz J. Meta-analysis: non-pathogenic yeast *Saccharomyces boulardii* in the prevention of antibiotic-associated diarrhoea. *Aliment Pharmacol Ther* 2005;22:365–72.
- [199] Bischoff SC, Escher J, Hebuterne X, Klek S, Krznaric Z, Schneider S, et al. ESPEN practical guideline: clinical nutrition in inflammatory bowel disease. *Clin Nutr* 2020;39:632–53.
- [200] Ng SC, Hart AL, Kamm MA, Stagg AJ, Knight SC. Mechanisms of action of probiotics: recent advances. *Inflamm Bowel Dis* 2009;15:300–10.
- [201] Morrow LE, Kollef MH. Probiotics in the intensive care unit: why controversies and confusion abound. *Crit Care* 2008;12:160.
- [202] Walker WA. Mechanisms of action of probiotics. *Clin Infect Dis* 2008;46(Suppl 2):S87–91. discussion S144–51.
- [203] Roberfroid M. Prebiotics: the concept revisited. *J Nutr* 2007;137: 830S–7S.
- [204] Chowdhury AH, Adiamah A, Kushairi A, Varadhan KK, Krznaric Z, Kulkarni AD, et al. Perioperative probiotics or synbiotics in adults undergoing elective abdominal surgery: a systematic review and meta-analysis of randomized controlled trials. *Ann Surg* 2019. <https://doi.org/10.1097/SLA.0000000000003581> [Epub ahead of print].
- [205] Besselink MG, van Santvoort HC, Buskens E, Boermeester MA, van Goor H, Timmerman HM, et al. Probiotic prophylaxis in predicted severe acute pancreatitis: a randomised, double-blind, placebo-controlled trial. *Lancet* 2008;371:651–9.
- [206] Gou S, Yang Z, Liu T, Wu H, Wang C. Use of probiotics in the treatment of severe acute pancreatitis: a systematic review and meta-analysis of randomized controlled trials. *Crit Care* 2014;18:R57.
- [207] Arvanitakis M, Ockenga J, Bezmarevic M, Gianotti L, Krznaric Z, Lobo DN, et al. ESPEN guideline on clinical nutrition in acute and chronic pancreatitis. *Clin Nutr* 2020;39:612–31.
- [208] Bischoff SC, Boirie Y, Cederholm T, Chourdakis M, Cuerda C, Delzenne NM, et al. Towards a multidisciplinary approach to understand and manage obesity and related diseases. *Clin Nutr* 2017;36:917–38.
- [209] Martin D, Roulin D, Grass F, Addor V, Ljungqvist O, Demartines N, et al. A multicentre qualitative study assessing implementation of an enhanced recovery after surgery program. *Clin Nutr* 2018;37:2172–7.